

**United States Department of the Interior
Bureau of Land Management**

**Environmental Assessment - Appendices
DOI-BLM-CO-N050-2020-0056**

***Piceance-East Douglas Herd Management Area
Gather and Fertility Control Plan***

November 2020

U.S. Department of the Interior
Bureau of Land Management
Northwest District
White River Field Office
220 East Market St
Meeker, CO 81641



BLM

Table of Contents

Appendix A. Maps	1
Appendix B. Review of Current Situation Memo	3
Appendix C. Excess Determination Memo	7
Appendix D. Animal Welfare Standards	9
Appendix E. Alternatives Considered but Not Analyzed in Detail	29
Appendix F. Population Model	34
Appendix G. Scientific Literature Review.....	34

APPENDIX A. MAPS

Figure 1. Piceance East-Douglas Herd Management Area





Figure 2. Lands with Wilderness Characteristics, Areas of Critical Environmental Concern, and National Historic District within PEDHMA



APPENDIX B. REVIEW OF CURRENT SITUATION MEMO



United States Department of the Interior

BUREAU OF LAND MANAGEMENT
White River Field Office
220 East Market Street, Meeker, Co. 81641
(970)878-3800




In Reply Refer To:
4720

October 19, 2020

Memorandum

To: Kent E. Walter, Field Manager, White River Field Office

From: Kyle Arnold, White River Field Office, Renewable Resources Staff Supervisor 

Subject: Removal of Excess Wild Horses in the Piceance-East Douglas Herd Management Area

Introduction

Wild horse populations within the White River Field Office (WRFO) are above management objectives as identified in the Resource Management Plan (and amendments) for the Piceance-East Douglas Herd Management Area (PEDHMA). Populations have continued to increase to a level that has never been reached in the WRFO. As the population increases inside the PEDHMA the occurrences of excess wild horses temporarily or permanently relocating outside of the PEDHMA increases due to the animals attempts to find the forage, water, cover, and space that they seek to survive. The population increase within the PEDHMA can be described as covering the entire boundary of the PEDHMA and at this time does not represent a thriving natural ecological balance of wild horses with other resource values in the WRFO. Due to the high numbers of wild horses located both inside and outside of the PEDHMA (levels that have never been reached before) the WRFO has and will continue to receive requests to remove excess wild horses from private lands with the most recent from August 2019 through August 2020.

Piceance-East Douglas Herd Management Area

Through previous Land Use Plan decisions the WRFO chose to manage wild horses in the PEDHMA, which was designated in 1986 for the long-term management of wild horses, on the selected range from the Piceance Basin Herd Unit, and the portion of the Douglas Creek Herd Unit east of Douglas Creek. In 1997, through the Resource Management Plan (RMP) the PEDHMA was expanded adding approximately 28,830 acres known as the Greasewood Allotment. The current PEDHMA is comprised of approximately 158,310 acres of public; 5,330 acres of state; and 26,490 acres of private lands totaling approximately 190,130 acres. The present day PEDHMA contained the largest concentration of wild horses during the original census conducted in 1974. 141 of the 154 wild horses counted during this inventory were found within the current PEDHMA boundary. In the 1980 Management Framework Plan, the Appropriate Management Level (AML) was set at 95-140 wild horses and would be managed within the selected range. this was carried forward in the 1997 RMP. The AML was adjusted upward in 2002 to the

INTERIOR REGION 7 • UPPER COLORADO BASIN
COLORADO, NEW MEXICO, UTAH, WYOMING

current range of 135-235 wild horses.

Rationale for a Gathering Excess Wild Horses

Private Land Requests

As noted above, the WRFO has and will continue to receive multiple requests from the private land owners both inside and outside of the PEDHMA to come gather excess wild horses due to the impacts associated with an overpopulation of wild horses described as: 1) consumption of meadows by wild horses results in not being able to utilize pasture because feed is removed; 2) fencing around the ranch does not keep the wild horses out; and 3) the wild horses are destroying water resources. Private land requests were one of the driving factors for the completion of the Tommys Draw fence in 2017 along with the Environmental Assessment to remove excess horses that had relocated outside of the PEDHMA.

The BLM's regulations at 43 CFR 4720.2-1 provide direction on removing excess wild horses when a request is received requesting removal of wild horses from private lands.

Wild Horse Population Estimates

In February 2016, the WRFO conducted a wild horse aerial inventory and that resulted in a count of 337 located within the PEDHMA. Since that inventory, the wild horse population has continued to increase at an approximate 20 percent recruitment rate that has been used for this region, bringing the current estimated PEDHMA population 838 wild horses after the 2020 foaling season. The Appropriate Management Level (AML) within the PEDHMA is 135-235 wild horses which results in a current estimated excess of between 603 to 703 wild horses just within the PEDHMA.

The WRFO has also identified excess wild horses that have relocated outside of the PEDHMA. Assuming a 20 percent recruitment rate each year since 2016, the estimated population of excess wild horses outside of the PEDHMA is at 438.

These current wild horse estimates within and outside the PEDHMA result in a total of 1,041 to 1,141 wild horses in excess within the WRFO.

Recommendation to Management

The WRFO must begin development of a gather plan to remove excess wild horses from within the PEDHMA (see attached map). As you know, the WRFO has designated the PEDHMA for the long-term management of wild horses. BLM Manual 4720 defines excess animals "as those animals which must be removed from an area to preserve and maintain a thriving natural ecological balance (TNEB) and multiple-use relationship in that area. This definition includes wild horses or burros located outside the HMA in areas not designated for their long-term maintenance" (BLM Manual 4720.12). It is important that the WRFO immediately begin development of a gather removal plan to address excess wild horses in


INTERIOR REGION 7 • UPPER COLORADO BASIN
COLORADO, NEW MEXICO, UTAH, WYOMING

accordance with 43 CFR 4720.2-1. NEPA is already in place to conduct wild horse gathers outside of the PEDHMA (doiblmcoN0520150023ea, doiblmcoN0520170056ea and doiblmcoN0520180071dna). Currently, staff is working on the NEPA document to remove excess wild horses located within the PEDHMA, it is on schedule to be completed this coming February 2021.

Staff recommendation is to remove excess wild horses impacting natural resources within the WRFO, within and outside of the PEDHMA. Wild horse population levels are resulting in overutilization of forage and water resources on public and private lands.

Attachment:
Map

INTERIOR REGION 7 • UPPER COLORADO BASIN
COLORADO, NEW MEXICO, UTAH, WYOMING



INTERIOR REGION 7 • UPPER COLORADO BASIN
COLORADO, NEW MEXICO, UTAH, WYOMING

APPENDIX C. EXCESS DETERMINATION MEMO



United States Department of the Interior

BUREAU OF LAND MANAGEMENT
White River Field Office
220 East Market Street
Meeker, CO 81641



In Reply Refer To: 4720
(CO-LLCON05000)

October 22, 2020

Memorandum

To: Catherine Cook, Acting Northwest District Manager

From: Kent E. Walter, Field Manager, White River Field Office

Subject: Removal of Excess Wild Horses from within the Piceance-East Douglas Herd Management Area

Introduction

Piceance-East Douglas Herd Management Area Wild Horse Population Estimates

In February 2016, the WRFO conducted a wild horse aerial inventory that resulted in a count of 337 wild horses located within the Piceance-East Douglas Herd Management Area (PEDHMA). Since that inventory, the wild horse population has continued to increase at an approximate 20 percent recruitment rate bringing the current estimated PEDHMA wild horse population to 838 wild horses by fall 2020. The Appropriate Management Level (AML) within the PEDHMA is 135-235 wild horses so at this time there is an estimated excess of between 603 to 703 wild horses just within the PEDHMA so currently at more than 3.5 times over the AML.

The WRFO has also identified excess wild horses that have relocated outside of the PEDHMA (in the North Piceance Herd Area, West Douglas Herd Area, and other areas beyond these boundaries). Continuing with a 20 percent recruitment rate each year since 2016 but less the gather operation of 2017 the current estimated population of excess wild horses outside of the PEDHMA is 438.

These excess wild horse estimates within and outside of the PEDHMA result in a current total of 1,041 to 1,141 inside the WRFO boundary. The current wild horse overpopulation levels are resulting in over utilization of the primary resources (forage and water) on public, state, and private lands within and outside the PEDHMA. As a result, this is negatively impacting our ability to persevere and maintain a thriving natural ecological balance (TNEB) and multiple-use relationship in that area.

Recommendation to Management

The WRFO has NEPA in place to conduct wild horses gathers outside of the PEDHMA (DOI-BLM-CO-N050-2015-0023-EA, DOI-BLM-CO-N050-2017-0056-EA, and DOI-BLM-CO-N050-

INTERIOR REGION 7 • UPPER COLORADO BASIN
COLORADO, NEW MEXICO, UTAH, WYOMING

2018-0071-DNA) but have lacked the approval to conducted such gather operations inside the PEDHMA.

The WRFO must begin development of a gather plan to remove excess wild horses from within the PEDHMA (see attached map). As you know, the PEDHMA was designated for the long-term management of wild horses. BLM Manual 4720 defines excess animals "... as those animals which must be removed from an area to preserve and maintain a thriving natural ecological balance (TNEB) and multiple-use relationship in that area. This definition includes wild horses and burros located outside of the HMA in areas not designated for their long-term maintenance." (BLM Manual 4720.12).

It is important that the WRFO immediately complete a 10-year gather and removal plan to address excess wild horses located within the PEDHMA in accordance with 43 CFR 4720.2-1. Currently, staff is working on that NEPA document to gather and remove excess wild horses and propose the initial and future use of fertility control treatments from within the PEDHMA.



INTERIOR REGION 7 • UPPER COLORADO BASIN
COLORADO, NEW MEXICO, UTAH, WYOMING

APPENDIX D. ANIMAL WELFARE STANDARDS

The following is Attachment 1 from Washington Office Instruction Memorandum 2015-151.

ATTACHMENT 1: COMPREHENSIVE ANIMAL WELFARE PROGRAM FOR WILD HORSE AND BURRO GATHERS

STANDARDS

Developed by

The Bureau of Land Management
Wild Horse and Burro Program

in collaboration with

Carolyn L. Stull, PhD
Kathryn E. Holcomb, PhD
University of California, Davis
School of Veterinary Medicine

June 30, 2015

June 30, 2015

CAWP Gather Standards

Attachment 1-1

WELFARE ASSESSMENT STANDARDS for GATHERS

CONTENTS

Welfare Assessment Standards

I. FACILITY DESIGN	2
A. Trap Site and Temporary Holding Facility	2
B. Loading and Unloading Areas.....	4
II. CAPTURE TECHNIQUE	5
A. Capture Techniques.....	5
B. Helicopter Drive Trapping	5
C. Roping	7
D. Bait Trapping.....	8
III. WILD HORSE AND BURRO CARE.....	8
A. Veterinarian.....	8
B. Care	9
C. Biosecurity	11
IV. HANDLING.....	12
A. Willful Acts of Abuse	12
B. General Handling	12
C. Handling Aids	12
V. TRANSPORTATION	13
A. General	13
B. Vehicles.....	14
C. Care of WH&Bs during Transport Procedures	15
VI. EUTHANASIA or DEATH.....	16
A. Euthanasia Procedures during Gather Operations.....	16
B. Carcass Disposal	17
Required documentation and responsibilities of Lead COR/COR/PI at gathers.....	18
Schematic of CAWP Gather Components.....	20

STANDARDS

Standard Definitions

Major Standard: Impacts the health or welfare of WH&Bs. Relates to an alterable equipment or facility standard or procedure. Appropriate wording is “must,” “unacceptable,” “prohibited.”

Minor Standard: unlikely to affect WH&Bs health or welfare or involves an uncontrollable situation. Appropriate wording is “should.”

Lead COR = Lead Contracting Officer’s Representative

COR = Contracting Officer’s Representative

PI = Project Inspector

WH&Bs = Wild horses and burros

I. FACILITY DESIGN

A. Trap Site and Temporary Holding Facility

1. The trap site and temporary holding facility must be constructed of stout materials and must be maintained in proper working condition, including gates that swing freely and latch or tie easily. **(major)**
2. The trap site should be moved close to WH&B locations whenever possible to minimize the distance the animals need to travel. **(minor)**
3. If jute is hung on the fence posts of an existing wire fence in the trap wing, the wire should be either be rolled up or let down for the entire length of the jute in such a way that minimizes the possibility of entanglement by WH&Bs unless otherwise approved by the Lead COR/COR/PI. **(minor)**
4. Fence panels in pens and alleys must be not less than 6 feet high for horses, 5 feet high for burros, and the bottom rail must not be more than 12 inches from ground level. **(major)**

5. The temporary holding facility must have a sufficient number of pens available to sort WH&Bs according to gender, age, number, temperament, or physical condition.
(**major**)
 - a. All pens must be assembled with capability for expansion. (**major**)
 - b. Alternate pens must be made available for the following: (**major**)
 - 1) WH&Bs that are weak or debilitated
 - 2) Mares/jennies with dependent foals
 - c. WH&Bs in pens at the temporary holding facility should be maintained at a proper stocking density such that when at rest all WH&Bs occupy no more than half the pen area. (minor)
6. An appropriate chute designed for restraining WH&Bs must be available for necessary procedures at the temporary holding facility. This does not apply to bait trapping operations unless directed by the Lead COR/COR/PI. (**major**)
7. There must be no holes, gaps or openings, protruding surfaces, or sharp edges present in fence panels or other structures that may cause escape or possible injury. (**major**)
8. Padding must be installed on the overhead bars of all gates and chutes used in single file alleys. (**major**)
9. Hinged, self-latching gates must be used in all pens and alleys except for entry gates into the trap, which may be secured with tie ropes. (**major**)
10. Finger gates (one-way funnel gates) used in bait trapping must be constructed of materials approved by the Lead COR/COR/PI. Finger gates must not be constructed of materials that have sharp ends that may cause injuries to WH&Bs, such as "T" posts, sharpened willows, etc. (**major**)
11. Water must be provided at a minimum rate of ten gallons per 1000 pound animal per day, adjusted accordingly for larger or smaller horses, burros and foals, and environmental conditions, with each trough placed in a separate location of the pen (i.e. troughs at opposite ends of the pen). Water must be refilled at least every morning and evening. (**major**)
12. The design of pens at the trap site and temporary holding facility should be constructed with rounded corners. (minor)

13. All gates and panels in the animal holding and handling pens and alleys of the trap site must be covered with materials such as plywood, snow fence, tarps, burlap, etc. approximately 48” in height to provide a visual barrier for the animals. All materials must be secured in place. **(major)**

These guidelines apply:

- a. For exterior fences, material covering panels and gates must extend from the top of the panel or gate toward the ground. **(major)**
 - b. For alleys and small internal handling pens, material covering panels and gates should extend from no more than 12 inches below the top of the panel or gate toward the ground to facilitate visibility of animals and the use of flags and paddles during sorting. (minor)
 - c. The initial capture pen may be left uncovered as necessary to encourage animals to enter the first pen of the trap. (minor)
14. Non-essential personnel and equipment must be located to minimize disturbance of WH&Bs. **(major)**
 15. Trash, debris, and reflective or noisy objects should be eliminated from the trap site and temporary holding facility. (minor)

B. Loading and Unloading Areas

1. Facilities in areas for loading and unloading WH&Bs at the trap site or temporary holding facility must be maintained in a safe and proper working condition, including gates that swing freely and latch or tie easily. **(major)**
2. The side panels of the loading chute must be a minimum of 6 feet high and fully covered with materials such as plywood or metal without holes that may cause injury. **(major)**
3. There must be no holes, gaps or openings, protruding surfaces, or sharp edges present in fence panels or other structures that may cause escape or possible injury. **(major)**
4. All gates and doors must open and close easily and latch securely. **(major)**

5. Loading and unloading ramps must have a non-slip surface and be maintained in a safe and proper working condition to prevent slips and falls. Examples of non-slip flooring would include, but not be limited to, rubber mats, sand, shavings, and steel reinforcement rods built into ramp. There must be no holes in the flooring or items that can cause an animal to trip. **(major)**
6. Trailers must be properly aligned with loading and unloading chutes and panels such that no gaps exist between the chute/panel and floor or sides of the trailer creating a situation where a WH&B could injure itself. **(major)**
7. Stock trailers should be positioned for loading or unloading such that there is no more than 12" clearance between the ground and floor of the trailer for burros and 18" for horses. **(minor)**

II. CAPTURE TECHNIQUE

A. Capture Techniques

1. WH&Bs gathered on a routine basis for removal or return to range must be captured by the following approved procedures under direction of the Lead COR/COR/PI. **(major)**
 - a. Helicopter
 - b. Bait trapping
2. WH&Bs must not be captured by snares or net gunning. **(major)**
3. Chemical immobilization must only be used for capture under exceptional circumstances and under the direct supervision of an on-site veterinarian experienced with the technique. **(major)**

B. Helicopter Drive Trapping

1. The helicopter must be operated using pressure and release methods to herd the animals in a desired direction and should not repeatedly evoke erratic behavior in the WH&Bs causing injury or exhaustion. Animals must not be pursued to a point of exhaustion; the on-site veterinarian must examine WH&Bs for signs of exhaustion. **(major)**

2. The rate of movement and distance the animals travel must not exceed limitations set by the Lead COR/COR/PI who will consider terrain, physical barriers, access limitations, weather, condition of the animals, urgency of the operation (animals facing drought, starvation, fire, etc.) and other factors. **(major)**
 - a. WH&Bs that are weak or debilitated must be identified by BLM staff or the contractors. Appropriate gather and handling methods should be used according to the direction of the Lead COR/COR/PI. **(major)**
 - b. The appropriate herding distance and rate of movement must be determined on a case-by-case basis considering the weakest or smallest animal in the group (e.g., foals, pregnant mares, or horses that are weakened by body condition, age, or poor health) and the range and environmental conditions present. **(major)**
 - c. Rate of movement and distance travelled must not result in exhaustion at the trap site, with the exception of animals requiring capture that have an existing severely compromised condition prior to gather. Where compromised animals cannot be left on the range or where doing so would only serve to prolong their suffering, euthanasia will be performed in accordance with BLM policy. **(major)**
3. WH&Bs must not be pursued repeatedly by the helicopter such that the rate of movement and distance travelled exceeds the limitation set by the Lead COR/COR/PI. Abandoning the pursuit or alternative capture methods may be considered by the Lead COR/COR/PI in these cases. **(major)**
4. When WH&Bs are herded through a fence line en route to the trap, the Lead COR/COR/PI must be notified by the contractor. The Lead COR/COR/PI must determine the appropriate width of the opening that the fence is let down to allow for safe passage through the opening. The Lead COR/COR/PI must decide if existing fence lines require marking to increase visibility to WH&Bs. **(major)**
5. The helicopter must not come into physical contact with any WH&B. The physical contact of any WH&B by helicopter must be documented by Lead COR/COR/PI along with the circumstances. **(major)**
6. WH&Bs may escape or evade the gather site while being moved by the helicopter. If there are mare/dependent foal pairs in a group being brought to a trap and half of an identified pair is thought to have evaded capture, multiple attempts by helicopter may

be used to bring the missing half of the pair to the trap or to facilitate capture by roping. In these instances, animal condition and fatigue must be evaluated by the Lead COR/COR/PI or on-site veterinarian on a case-by-case basis to determine the number of attempts that can be made to capture an animal. **(major)**

7. Horse captures must not be conducted when ambient temperature at the trap site is below 10°F or above 95°F without approval of the Lead COR/COR/PI. Burro captures must not be conducted when ambient temperature is below 10°F or above 100°F without approval of the Lead COR/COR/PI. The Lead COR/COR/PI will not approve captures when the ambient temperature exceeds 105 °F. **(major)**

C. Roping

1. The roping of any WH&B must be approved prior to the procedure by the Lead COR/COR/PI. **(major)**
2. The roping of any WH&B must be documented by the Lead COR/COR/PI along with the circumstances. WH&Bs may be roped under circumstances which include but are not limited to the following: reunite a mare or jenny and her dependent foal; capture nuisance, injured or sick WH&Bs or those that require euthanasia; environmental reasons such as deep snow or traps that cannot be set up due to location or environmentally sensitive designation; and public and animal safety or legal mandates for removal. **(major)**
3. Ropers should dally the rope to their saddle horn such that animals can be brought to a stop as slowly as possible and must not tie the rope hard and fast to the saddle so as to intentionally jerk animals off their feet. **(major)**
4. WH&Bs that are roped and tied down in recumbency must be continuously observed and monitored by an attendant at a maximum of 100 feet from the animal. **(major)**
5. WH&Bs that are roped and tied down in recumbency must be untied within 30 minutes. **(major)**
6. If the animal is tied down within the wings of the trap, helicopter drive trapping within the wings will cease until the tied-down animal is removed. **(major)**
7. Sleds, slide boards, or slip sheets must be placed underneath the animal's body to move and/or load recumbent WH&Bs. **(major)**

8. Halters and ropes tied to a WH&B may be used to roll, turn, position or load a recumbent animal, but a WH&B must not be dragged across the ground by a halter or rope attached to its body while in a recumbent position. **(major)**
9. Animals captured by roping must be evaluated by the on-site/on-call veterinarian within four hours after capture, marked for identification at the trap site, and be re-evaluated periodically as deemed necessary by the on-site/on-call veterinarian. **(major)**

D. Bait Trapping

1. WH&Bs may be lured into a temporary trap using bait (feed, mineral supplement, water) or sexual attractants (mares/jennies in heat) with the following requirements:
 - a. The period of time water sources other than in the trap site are inaccessible must not adversely affect the wellbeing of WH&Bs, wildlife or livestock, as determined by the Lead COR/COR/PI. **(major)**
 - b. Unattended traps must not be left unobserved for more than 12 hours. **(major)**
 - c. Mares/jennies and their dependent foals must not be separated unless for safe transport. **(major)**
 - d. WH&Bs held for more than 12 hours must be provided with accessible clean water at a minimum rate of ten gallons per 1000 pound animal per day, adjusted accordingly for larger or smaller horses, burros and foals and environmental conditions. **(major)**
 - e. WH&Bs held for more than 12 hours must be provided good quality hay at a minimum rate of 20 pounds per 1000 pound adult animal per day, adjusted accordingly for larger or smaller horses, burros and foals. **(major)**
 - 1) Hay must not contain poisonous weeds, debris, or toxic substances. **(major)**
 - 2) Hay placement must allow all WH&Bs to eat simultaneously. **(major)**

III. WILD HORSE AND BURRO CARE

A. Veterinarian

1. On-site veterinary support must be provided for all helicopter gathers and on-site or on-call support must be provided for bait trapping. **(major)**

2. Veterinary support must be under the direction of the Lead COR/COR/PI. The on-site/on-call veterinarian will provide consultation on matters related to WH&B health, handling, welfare, and euthanasia at the request of the Lead COR/COR/PI. All decisions regarding medical treatment or euthanasia will be made by the on-site Lead COR/COR/PI. **(major)**

B. Care

1. Feeding and Watering
 - a. Adult WH&Bs held in traps or temporary holding pens for longer than 12 hours must be fed every morning and evening with water available at all times other than when animals are being sorted or worked. **(major)**
 - b. Water must be provided at a minimum rate of ten gallons per 1000 pound animal per day, adjusted accordingly for larger or smaller horses, burros and foals, and environmental conditions, with each trough placed in a separate location of the pen (i.e. troughs at opposite ends of the pen). **(major)**
 - c. Good quality hay must be fed at a minimum rate of 20 pounds per 1000 pound adult animal per day, adjusted accordingly for larger or smaller horses, burros and foals. **(major)**
 - i. Hay must not contain poisonous weeds or toxic substances. **(major)**
 - ii. Hay placement must allow all WH&Bs to eat simultaneously. **(major)**
 - d. When water or feed deprivation conditions exist on the range prior to the gather, the Lead COR/COR/PI should adjust the watering and feeding arrangements in consultation with the onsite veterinarian as necessary to provide for the needs of the animals. **(minor)**
2. Dust abatement
 - a. Dust abatement by spraying the ground with water must be employed when necessary at the trap site and temporary holding facility. **(major)**

3. Trap Site

- a. Dependent foals or weak/debilitated animals must be separated from other WH&Bs at the trap site to avoid injuries during transportation to the temporary holding facility. Separation of dependent foals from mares must not exceed four hours unless the Lead COR/COR/PI authorizes a longer time or a decision is made to wean the foals. **(major)**

4. Temporary Holding Facility

- a. All WH&Bs in confinement must be observed at least once daily to identify sick or injured WH&Bs and ensure adequate food and water. **(major)**
- b. Foals must be reunited with their mares/jennies at the temporary holding facility within four hours of capture unless the Lead COR/COR/PI authorizes a longer time or foals are old enough to be weaned during the gather. **(major)**
- c. Non-ambulatory WH&Bs must be located in a pen separate from the general population and must be examined by the BLM horse specialist and/or on-call or on-site veterinarian as soon as possible, no more than four hours after recumbency is observed. Unless otherwise directed by a veterinarian, hay and water must be accessible to an animal within six hours after recumbency. **(major)**
- d. Alternate pens must be made available for the following: **(major)**
 - 1) WH&Bs that are weak or debilitated
 - 2) Mares/jennies with dependent foals
- e. Aggressive WH&Bs causing serious injury to other animals should be identified and relocated into alternate pens when possible. (minor)
- f. WH&Bs in pens at the temporary holding facility should be maintained at a proper stocking density such that when at rest all WH&Bs occupy no more than half the pen area. (minor)

C. Biosecurity

1. Health records for all saddle and pilot horses used on WH&B gathers must be provided to the Lead COR/COR/PI prior to joining a gather, including: **(major)**
 - a. Certificate of Veterinary Inspection (Health Certificate, within 30 days).
 - b. Proof of:
 - 1) A negative test for equine infectious anemia (Coggins or EIA ELISA test) within 12 months.
 - 2) Vaccination for tetanus, eastern and western equine encephalomyelitis, West Nile virus, equine herpes virus, influenza, *Streptococcus equi*, and rabies within 12 months.
2. Saddle horses, pilot horses and mares used for bait trapping lures must not be removed from the gather operation (such as for an equestrian event) and allowed to return unless they have been observed to be free from signs of infectious disease for a period of at least three weeks and a new Certificate of Veterinary Examination is obtained after three weeks and prior to returning to the gather. **(major)**
3. WH&Bs, saddle horses, and pilot horses showing signs of infectious disease must be examined by the on-site/on-call veterinarian. **(major)**
 - a. Any saddle or pilot horses showing signs of infectious disease (fever, nasal discharge, or illness) must be removed from service and isolated from other animals on the gather until such time as the horse is free from signs of infectious disease and approved by the on-site/on-call veterinarian to return to the gather. **(major)**
 - b. Groups of WH&Bs showing signs of infectious disease should not be mixed with groups of healthy WH&Bs at the temporary holding facility, or during transport. **(minor)**
4. Horses not involved with gather operations should remain at least 300 yards from WH&Bs, saddle horses, and pilot horses being actively used on a gather. **(minor)**

IV. HANDLING

A. Willful Acts of Abuse

1. Hitting, kicking, striking, or beating any WH&B in an abusive manner is prohibited. **(major)**
2. Dragging a recumbent WH&B without a sled, slide board or slip sheet is prohibited. Ropes used for moving the recumbent animal must be attached to the sled, slide board or slip sheet unless being loaded as specified in Section II. C. 8. **(major)**
3. There should be no deliberate driving of WH&Bs into other animals, closed gates, panels, or other equipment. (minor)
4. There should be no deliberate slamming of gates and doors on WH&Bs. (minor)
5. There should be no excessive noise (e.g., constant yelling) or sudden activity causing WH&Bs to become unnecessarily flighty, disturbed or agitated. (minor)

B. General Handling

1. All sorting, loading or unloading of WH&Bs during gathers must be performed during daylight hours except when unforeseen circumstances develop and the Lead COR/CO/PI approves the use of supplemental light. **(major)**
2. WH&Bs should be handled to enter runways or chutes in a forward direction. (minor)
3. WH&Bs should not remain in single-file alleyways, runways, or chutes longer than 30 minutes. (minor)
4. Equipment except for helicopters should be operated and located in a manner to minimize flighty behavior. (minor)

C. Handling Aids

1. Handling aids such as flags and shaker paddles must be the primary tools for driving and moving WH&Bs during handling and transport procedures. Contact of the flag or paddle end of primary handling aids with a WH&B is allowed. Ropes looped around the hindquarters may be used from horseback or on foot to assist in moving an animal forward or during loading. **(major)**

2. Electric prods must not be used routinely as a driving aid or handling tool. Electric prods may be used in limited circumstances only if the following guidelines are followed:
 - a. Electric prods must only be a commercially available make and model that uses DC battery power and batteries should be fully charged at all times. **(major)**
 - b. The electric prod device must never be disguised or concealed. **(major)**
 - c. Electric prods must only be used after three attempts using other handling aids (flag, shaker paddle, voice or body position) have been tried unsuccessfully to move the WH&Bs. **(major)**
 - d. Electric prods must only be picked up when intended to deliver a stimulus; these devices must not be constantly carried by the handlers. **(major)**
 - e. Space in front of an animal must be available to move the WH&B forward prior to application of the electric prod. **(major)**
 - f. Electric prods must never be applied to the face, genitals, anus, or underside of the tail of a WH&B. **(major)**
 - g. Electric prods must not be applied to any one WH&B more than three times during a procedure (e.g., sorting, loading) except in extreme cases with approval of the Lead COR/COR/PI. Each exception must be approved at the time by the Lead COR/COR/PI. **(major)**
 - h. Any electric prod use that may be necessary must be documented daily by the Lead COR/COR/PI including time of day, circumstances, handler, location (trap site or temporary holding facility), and any injuries (to WH&B or human). **(major)**

V. TRANSPORTATION

A. General

1. All sorting, loading, or unloading of WH&Bs during gathers must be performed during daylight hours except when unforeseen circumstances develop and the Lead COR/CO/PI approves the use of supplemental light. **(major)**

2. WH&Bs identified for removal should be shipped from the temporary holding facility to a BLM facility within 48 hours. (minor)
 - a. Shipping delays for animals that are being held for release to range or potential on-site adoption must be approved by the Lead COR/COR/PI. (**major**)
3. Shipping should occur in the following order of priority; 1) debilitated animals, 2) pairs, 3) weanlings, 4) dry mares and 5) studs. (minor)
4. Planned
5. transport time to the BLM preparation facility from the trap site or temporary holding facility must not exceed 10 hours. (**major**)
6. WH&Bs should not wait in stock trailers and/or semi-trailers at a standstill for more than a combined period of three hours during the entire journey. (minor)

B. Vehicles

1. Straight-deck trailers and stock trailers must be used for transporting WH&Bs. (**major**)
 - a. Two-tiered or double deck trailers are prohibited. (**major**)
 - b. Transport vehicles for WH&Bs must have a covered roof or overhead bars containing them such that WH&Bs cannot escape. (**major**)
2. WH&Bs must have adequate headroom during loading and unloading and must be able to maintain a normal posture with all four feet on the floor during transport without contacting the roof or overhead bars. (**major**)
3. The width and height of all gates and doors must allow WH&Bs to move through freely. (**major**)
4. All gates and doors must open and close easily and be able to be secured in a closed position. (**major**)
5. The rear door(s) of the trailers must be capable of opening the full width of the trailer. (**major**)
6. Loading and unloading ramps must have a non-slip surface and be maintained in proper working condition to prevent slips and falls. (**major**)

7. Transport vehicles more than 18 feet and less than 40 feet in length must have a minimum of one partition gate providing two compartments; transport vehicles 40 feet or longer must have at least two partition gates to provide a minimum of three compartments. **(major)**
8. All partitions and panels inside of trailers must be free of sharp edges or holes that could cause injury to WH&Bs. **(major)**
9. The inner lining of all trailers must be strong enough to withstand failure by kicking that would lead to injuries. **(major)**
10. Partition gates in transport vehicles should be used to distribute the load into compartments during travel. **(minor)**
11. Surfaces and floors of trailers must be cleaned of dirt, manure and other organic matter prior to the beginning of a gather. **(major)**

C. Care of WH&Bs during Transport Procedures

1. WH&Bs that are loaded and transported from the temporary holding facility to the BLM preparation facility must be fit to endure travel. **(major)**
 - a. WH&Bs that are non-ambulatory, blind in both eyes, or severely injured must not be loaded and shipped unless it is to receive immediate veterinary care or euthanasia. **(major)**
 - b. WH&Bs that are weak or debilitated must not be transported without approval of the Lead COR/COR/PI in consultation with the on-site veterinarian. Appropriate actions for their care during transport must be taken according to direction of the Lead COR/COR/PI. **(major)**
2. WH&Bs should be sorted prior to transport to ensure compatibility and minimize aggressive behavior that may cause injury. **(minor)**
3. Trailers must be loaded using the minimum space allowance in all compartments as follows: **(major)**
 - a. 12 square feet per adult horse.
 - b. 6.0 square feet per dependent horse foal.
 - c. 8.0 square feet per adult burro.
 - d. 4.0 square feet per dependent burro foal.

4. The Lead COR/COR/PI in consultation with the receiving Facility Manager must document any WH&B that is recumbent or dead upon arrival at the destination.
(major)
 - a. Non-ambulatory or recumbent WH&Bs must be evaluated on the trailer and either euthanized or removed from the trailers using a sled, slide board or slip sheet.
(major)
5. Saddle horses must not be transported in the same compartment with WH&Bs.
(major)

VI. EUTHANASIA OR DEATH

A. Euthanasia Procedure during Gather Operations

1. An authorized, properly trained, and experienced person as well as a firearm appropriate for the circumstances must be available at all times during gather operations. When the travel time between the trap site and temporary holding facility exceeds one hour or if radio or cellular communication is not reliable, provisions for euthanasia must be in place at both the trap site and temporary holding facility during the gather operation. (major)
2. Euthanasia must be performed according to American Veterinary Medical Association euthanasia guidelines (2013) using methods of gunshot or injection of an approved euthanasia agent. (major)
3. The decision to euthanize and method of euthanasia must be directed by the Authorized Officer or their Authorized Representative(s) that include but are not limited to the Lead COR/COR/PI who must be on site and may consult with the on-site/on-call veterinarian. (major)
4. Photos needed to document an animal's condition should be taken prior to the animal being euthanized. No photos of animals that have been euthanized should be taken. An exception is when a veterinarian or the Lead COR/COR/PI may want to document certain findings discovered during a postmortem examination or necropsy. (minor)
5. Any WH&B that dies or is euthanized must be documented by the Lead COR/COR/PI including time of day, circumstances, euthanasia method, location, a

description of the age, gender, and color of the animal and the reason the animal was euthanized. **(major)**

6. The on-site/on-call veterinarian should review the history and conduct a postmortem physical examination of any WH&B that dies or is euthanized during the gather operation. A necropsy should be performed whenever feasible if the cause of death is unknown. (minor)

B. Carcass Disposal

1. The Lead COR/COR/PI must ensure that appropriate equipment is available for the timely disposal of carcasses when necessary on the range, at the trap site, and temporary holding facility. **(major)**
2. Disposal of carcasses must be in accordance with state and local laws. **(major)**
3. WH&Bs euthanized with a barbiturate euthanasia agent must be buried or otherwise disposed of properly. **(major)**
4. Carcasses left on the range should not be placed in washes or riparian areas where future runoff may carry debris into ponds or waterways. Trenches or holes for buried animals should be dug so the bottom of the hole is at least 6 feet above the water table and 4-6 feet of level earth covers the top of the carcass with additional dirt mounded on top where possible. (minor)

CAWP
REQUIRED DOCUMENTATION AND RESPONSIBILITIES OF LEAD
COR/COR/PI

Required Documentation

Section	Documentation
II.B.5	Helicopter contact with any WH&B.
II.C.2	Roping of any WH&B.
III.B.3.a and	Reason for allowing longer than four hours to reunite foals with mares/jennies. Does not apply if foals are being weaned.
III.B.4.b	
III.C.1	Health status of all saddle and pilot horses.
IV.C.2.h	All uses of electric prod.
V.C.4	Any WH&B that is recumbent or dead upon arrival at destination following transport.
VI.A.5	Any WH&B that dies or is euthanized during gather operation.

Responsibilities

Section	Responsibility
I.A.10	Approve materials used in construction of finger gates in bait trapping
II.A.1	Direct gather procedures using approved gather technique.
II.B.2	Determine rate of movement and distance limitations for WH&B helicopter gather.
II.B.2.a	Direct appropriate gather/handling methods for weak or debilitated WH&B.
II.B.3	Determine whether to abandon pursuit or use other capture method in order to avoid repeated pursuit of WH&B.
II.B.4	Determine width and need for visibility marking when using opening in fence en route to trap.
II.B.6	Determine number of attempts that can be made to capture the missing half of a mare/foal pair that has become separated.
II.B.7	Determine whether to proceed with gather when ambient temperature is outside the range of 10°F to 95°F for horses or 10°F to 100°F for burros.
II.C.1	Approve roping of any WH&B.
II.D.1.a	Determine period of time that water outside a bait trap is inaccessible such that wellbeing of WH&Bs, wildlife, or livestock is not adversely affected.
III.A.2	Direct and consult with on-site/on-call veterinarian on any matters related to WH&B health, handling, welfare and euthanasia.

- III.B.1.e Adjust feed/water as necessary, in consultation with onsite/on call veterinarian, to provide for needs of animals when water or feed deprivation conditions exist on range.
- III.B.4.c Determine provision of water and hay to non-ambulatory animals.
- IV.C.2.g Approve use of electric prod more than three times, for exceptional cases only.
- V.A.1 Approve sorting, loading, or unloading at night with use of supplemental light.
- V.A.2.a Approve shipping delays of greater than 48 hours from temporary holding facility to BLM facility.
- V.C.1.b Approve of transport and care during transport for weak or debilitated WH&B.
- VI.A.3 Direct decision regarding euthanasia and method of euthanasia for any WH&B; may consult with on-site/on-call veterinarian.
- VI.B.1 Ensure that appropriate equipment is available for carcass disposal.

APPENDIX E. ALTERNATIVES CONSIDERED BUT NOT ANALYZED IN DETAIL

The following alternatives were considered but dismissed from detailed analysis for the reasons described below.

E.1. Alternative Gather Methods

Bait Trap Only

The BLM considered the use of bait-trapping as the primary gathering method. The use of bait-trapping, though effective in specific areas and circumstances, would be time-consuming and not cost-effective, considering the historical difficulties of gathering excess wild horses within the Piceance-East Douglas Herd Management Area (PEDHMA). Bait-trapping would also be impractical as the primary gather method for the PEDHMA due to the number of excess wild horses that need to be gathered and removed. However, bait-trapping could be used to help achieve Appropriate Management Level (AML) if gather efficiencies are too low using a helicopter or if a helicopter gather cannot be scheduled. This alternative was dismissed from detailed analysis as the primary gather method for the following reasons: 1) the project area is too large to effectively use this gather method; 2) road access for vehicles not associated with the gather to potential trapping locations is difficult to restrict and reduces gather efficiency; 3) road access necessary to get equipment in/out as well as safely transport gathered wild horses is limited; the amount of time required to come close to management objectives, and 4) the presence of scattered water resources on state, private and public lands inside the PEDHMA (some with no access) would make it almost impossible to restrict wild horse access to the extent necessary to gather and remove the excess animals if using bait-trapping achieve management goals.

Use of Alternative Capture Techniques

In other field offices, the public has suggested using capture methods (other than helicopters or bait-trapping) to gather excess wild horses. As no specific alternative methods were suggested, the BLM identified chemical immobilization, net-gunning, and wrangler/horseback drive trapping as potential methods for gathering wild horses. Net-gunning techniques normally used to capture big game also rely on helicopters. Chemical immobilization is a very specialized technique and strictly regulated. Currently the BLM does not have sufficient expertise to implement either of these methods and they would be impractical to use given the size of the PEDHMA, access limitations, number to be removed and approachability of the wild horses.

Use of wranglers on horseback drive-trapping to remove excess wild horses can be fairly effective on a small scale; but due to the number of excess wild horses to be removed, the large geographic size of the PEDHMA, access limitations, and approachability of the wild horses this technique would be ineffective and impractical. Horseback drive-trapping is also very labor intensive and can be harmful to the domestic horses and the wranglers used to herd the wild horses. For these reasons, this alternative was eliminated from further consideration for a large-scale gather.

E.2. Alternative Fertility Control Options

Using Field Darting to Deliver Fertility Treatments to Reduce Total Population Over Time

The BLM considered administering PZP or Gonacon in the one-year dose inoculations by field darting the mares. These methods are currently approved for use and is being used by BLM in other HMAs. This alternative (darting fertility control only, no removals) was dismissed from detailed analysis because the reduction in population growth is not sufficient to reduce overutilization of rangeland resources by wild horses and thus would not meet the purpose and need of the action. For these reasons, this alternative was determined to not be an effective or feasible method for managing wild horse populations and reducing rangeland impacts from over population in a timely manner within the PEDHMA.

Using Bait Trapping to Deliver Fertility Treatments

Population modeling (Appendix F) was completed to analyze the potential impacts associated with conducting gathers about every 2-3 years over the next 20-year period to treat captured mares with fertility control. Under this alternative, no excess wild horses would be removed. While the average population growth would be reduced, AML would not be achieved and the damage to the range associated with wild horse overpopulation would continue. This alternative would not meet the Purpose and Need for the Action was dismissed from further study.

Gather and Release All (including Excess) Wild Horses Every Two Years and Apply Two Year PZP or Other Contraceptive Vaccine to Wild Horses For Release

Another alternative to gather a substantial portion of the existing population (90 percent) and implement fertility control treatment only, without removal of excess wild horses using a two-year gather/treatment interval over a 10-year period, based on expected effectiveness of PZP-22 pellet vaccine or similar (e.g., GonaCon). The wild horse population would continue to have an average population growth rate of 2.3 to 13.7 percent, thus adding to the current wild horse overpopulation, albeit at a slower rate of growth than the No Action Alternative. The WinEquus modeling showed that implementing this kind of strategy would not eliminate continued wild horse management in the PEDHMA utilizing this alternative. This alternative would not decrease the existing overpopulation of wild horses, resource concerns and rangeland deterioration would continue, and implementation would result in substantially increased gather and fertility control costs relative to the alternatives that remove excess wild horses to the range prescribed by AML. In addition to not achieving AML, the time needed to complete a gather would also increase over time, because the more frequently an area is gathered, the gathering of wild horses becomes increasingly more difficult. Wild horses learn to evade the helicopter by taking cover in densely treed areas and access other areas generally in accessible (e.g. canyons). Wild horses tend to move out of an area when they hear a helicopter thereby further reducing the overall gather efficiency. Frequent gathers would increase the number of times wild horses would experience stress as individuals and the entire band/herd. It would become increasingly more difficult over time to repeat gather operations every two years to successfully treat a large portion of the population. For these reasons, this alternative was dropped from detailed study.

E.3. Alternatives Related to Population Size or Structure

Provide Supplemental Feed and Water

To accommodate higher populations of wild horses it has been suggested during public questions and comments that the BLM provide supplemental feed (hay) and/or haul water other than during a short-term emergency situation. Such an approach would not meet the definition of minimum feasible management and is inconsistent with current law, regulation, and policy (refer to 43 CFR 4710.4).

Return a Portion of the Population as a Non-Breeding Population

This alternative would involve capturing, gelding and/or spaying, and returning a portion of the population as a non-breeding population after the population is brought to the low end of AML. This alternative was not brought forward for detailed analysis because it is inconsistent with the 1997 RMP's direction (page 2-26) to manage wild horses in the PEDHMA "to provide a healthy, viable breeding population with a diverse age structure."

Utilize Only Sex Ratio Adjustment to Reduce Population Recruitment

This alternative would involve gathering and releasing all wild horses back into the PEDHMA under a sex ratio adjustment where more males than females would exist in order to bring the current estimated 20% recruitment rate down in order to curb gather operation occurrences. This alternative would be labor intensive and require additional years to reach a reduced recruitment rate. For those reasons, this alternative did not receive further consideration.

Gather to the AML Upper Limit

A post-gather population size at the upper level of the AML range would result in the AML being exceeded with the next foaling season. This would be unacceptable for several reasons as follows.

The AML represents "that 'optimum number' of wild horses that results in a thriving natural ecological balance and avoids a deterioration of the range" (Animal Protection Institute, 109 Interior Board of Land Appeals (IBLA) 119; 1989). The IBLA has also held that, "Proper range management dictates removal of horses before the herd size causes damage to the rangeland. Thus, the optimum number of horses is somewhere below the number that would cause resource damage." Refer to Animal Protection Institute, 118 IBLA 63, 75; 1991.

The upper level of the AML established for the PEDHMA represents the maximum population for which thriving natural ecological balance would be maintained. The lower level represents the number of animals to remain in the PEDHMA following a wild horse gather, in order to allow for a periodic gather cycle, and to prevent the population from exceeding the established AML between gathers.

Additionally, gathering to the upper range of AML would result in the need to follow up with another gather within one year (with resulting stress on the wild horse population) reducing gather efficiency overtime, and could result in overutilization of vegetation resources and damage to the rangeland if the BLM is unable to gather the excess wild horses in the PEDHMA on an annual basis. This alternative would not reduce the wild horse population growth rate of 20 percent in the PEDHMA and the BLM would not be able to conduct periodic gathers and still

maintain a thriving natural ecological balance. For these reasons, this alternative did not receive further consideration in this document.

Wild Horse Numbers Controlled by Natural Means

This alternative was eliminated from further consideration because it is contrary to the Wild and Free-Roaming Horse and Burro Act of 1971 (WFRHBA) that requires the BLM to prevent the range from deterioration associated with an overpopulation of wild horses. The alternative of using natural controls to achieve a desirable AML has not been shown to be feasible in the past. Wild horses in the PEDHMA are not substantially regulated by predators (which includes mountain lions and bears). In addition, wild horses are a long-lived species with documented foal survival rates exceeding 95 percent and they are not a self-regulating species. This alternative would result in a steady increase in numbers which would continually exceed the carrying capacity of the range until severe and unusual conditions that occur periodically -- such as blizzards or extreme drought -- cause catastrophic mortality of wild horses.

E.4. Alternatives Inconsistent with Existing Land Use Plan Allocations

Return the PEDHMA To Herd Area Status with Zero AML

Another alternative which has been suggested is to return the PEDHMA to Herd Area status and establish the AML as “0” animals due to resource concerns such as lack of forage, lack of water, and conflicts with other resources. The PEDHMA is expected to provide the necessary forage, water resources, cover and space to support a wild horse population of 135-235 animals. Additionally, this alternative would not be in conformance with existing land use plan decisions in the 1997 RMP because it is identified as an HMA rather than an HA.

Manage the Entire Population as a Non-Breeding Population

A management alternative which has been suggested is to manage the PEDHMA in its entirety as a non-breeding population of wild horses (e.g. entire herd of geldings). This alternative is not in conformance with the direction in the 1997 RMP, as amended to manage wild horses in the PEDHMA “to provide a healthy, viable breeding population with a diverse age structure” (page 2-26).

Remove Livestock within the PEDHMA

This alternative would involve no removal of wild horses and instead address the wild horse numbers and associated consumption of forage, through the removal of livestock within the PEDHMA. This alternative was not brought forward for detailed analysis because it is not in conformance with the 1997 RMP which allows for livestock grazing within the PEDHMA; the elimination of livestock grazing in an area would require an amendment to the RMP.

In previously analyzed the DOI-BLM-CO-110-2011-0058-EA, the BLM considered incrementally reducing forage allocated to livestock in the PEDHMA as wild horse populations continued to increase without intervention from a gather (Alternative C). In section 4.6.4 of the 2011 EA, the analysis demonstrated that the unchecked population growth of wild horses would within a few years eliminate forage allocations for livestock. Effectively eliminating forage allocations through an implementation decision (i.e., grazing permit) would be essentially similar

to a decision to remove livestock from the PEDHMA, which, as described above, is inconsistent with the 1997 RMP.

APPENDIX F. POPULATION MODEL

F.1. Population Model Overview

Population modeling is a tool designed to help BLM evaluate various management alternatives and possible outcomes for management of wild horses. The population model is not applicable for burros.

The WinEquus program, developed by Dr. Steven Jenkins at the University of Nevada at Reno was designed to assist wild horse and burro specialists in evaluating various management alternatives that could be considered for a particular area.

The model uses data on average survival probabilities and foaling rates of horses to simulate population growth for up to 20 years. The model accounts for year-to-year variation in these demographic parameters by using a randomization process to select survival probabilities and foaling rates for each age class from a distribution of values based on these averages. This aspect of population dynamics is called environmental stochasticity and reflects the fact that future environmental conditions that may affect horse populations cannot be known in advance. Therefore, each trial within the model will give a different pattern of population growth. Some trials may include mostly “good years”, when the population grows rapidly; other trials may include a series of several “bad” years in succession. The stochastic approach to population modeling uses repeated trials to project a range of possible population trajectories over a period of years, which is more realistic than predicting a single specific trajectory.

The model can incorporate selective removal and fertility control treatment as management strategies. A simulation may include no management, selective removal, fertility control treatment, or both removal and fertility control treatment. BLM can specify many different options for these management strategies such as the schedule of gathers for removal or fertility control treatment, the threshold population size which triggers a gather, the target population size following a removal, the ages and sexes of horses to be removed, and the effectiveness of fertility control treatment.

Modeling was completed for the Piceance-East Douglas Herd Management Area (PEDHMA), where BLM expects to only be able to gather 80 percent of the wild horses during gather operations. Population modeling was completed for all alternatives including the No Action. Initial population age structures that were utilized computed a stable age-sex distribution with the initial population size of 838 (the current estimated population of the PEDHMA). The percentages for females/males in the stable population used in the model were approximately 46 percent females and 54 percent males from the Garfield Flat HMA. All simulations used the survival probabilities and foaling rates supplied with the WinEquus population model for the Garfield Flat HMA. Survival data was collected by M. Ashley and S. Jenkins at Garfield Flat, Nevada between 1993 and 1999. Marked individuals were followed for a total of 708 animal-years to generate these survival probabilities. Foaling rate data was collected by M. Ashley and S. Jenkins at Garfield Flat, Nevada between 1993 and 1999. Marked females were followed for a total of 351 animal-years to generate these data on foaling rates.

These initial populations for the PEDHMA were entered into the model and put through simulations that included Gather to Low End of AML and Fertility Control Treatment (Alternative A), Gather to Low End of AML and Do Not Use Fertility Control Treatments (Alternative B), and No Action Alternative (Alternative C). Gate cut was also used to represent removing as many animals as possible from the PEDHMA to reach AML. The gate cut component does not represent what may occur on the ground as selective removal criteria may be utilized to reach AML to retain desirable characteristics in the PEDHMA over the long term. One additional difficulty with the model is that it does not have the ability to estimate the effectiveness of GonaCon as the preferred fertility control treatment. GonaCon could have longer lasting fertility control within the population which could increase the time that the PEDHMA stays within the AML. The simulations were run for 100 trials for the eleven years. For each simulation, a series of graphs and tables were provided which included the “most typical” trial, population sizes, growth rates, and gather numbers.

F.2. Results of Population Modeling

Out of the 100 trials in each simulation run, the model tabulated minimum, average, and maximum population sizes. The model was run for a period of eleven years from 2020 to 2030 and gives output through 2030. These numbers are useful to make relative comparisons of the different alternatives, and potential outcomes under different management options. The lowest, median and highest trials are displayed for each simulation completed. This output, together with the time series and most typical trial graphs are useful representations of the results of the program in terms of assessing the effects of the management alternatives because it shows not only expected average results but also extreme results that might be possible. The minimum population size in general reflects the numbers that would remain following management or random environmental impacts. The maximum population size generally reflects the population that existed prior to the gather, and in many cases that figure would not be exceeded during the ten years of the simulations. Half of the trials were greater than the median and half of them less than the median.

Under Table 1 and 2 below, the model depicts in population numbers that there is only a marginal difference, but what it does not show is that with gather operations and the use of fertility control treatments that the real difference, besides the frequency of gather operations necessary to stay within AML, is that a lower number of wild horses would need to be gathered and removed in order to stay within AML. So overall there is a benefit to the successful utilization of fertility control treatments in the PEDHMA.

Table 1. Population Size – Gather to Low End of AML and Fertility Control Treatment (Alternative A)

Estimated Population Sizes in 11 Years			
Trial	Minimum	Average	Maximum
Lowest	120	263	843
Median	152	284	906
Highest	175	333	1,145

Table 2. Population Size – Gather to Low End of AML and Do Not Use Fertility Control Treatments (Alternative B)

Estimated Population Sizes in 11 Years			
Trial	Minimum	Average	Maximum
Lowest	121	260	843
Median	155	284	912
Highest	182	339	1,163

Table 3. Population Size – No Action Alternative (Alternative C)


*Estimated Population Sizes in 10 Years versus 11 Years			
Trial	Minimum	Average	Maximum
Lowest	842	1,698	3,124
Median	924	2,635	5,586
Highest	1,176	3,503	8,311

F.2.1. Time Series Graph of Most Typical Trial

Based on the results from the model, spaghetti graphs (see below) were generated for each simulation. These graphs show how population size changes over time. The Y-axis scale remains constant for each graph; however, the X-axis was determined based on results and was unable to be changed. At first glance, there appears to be not much difference between the trials, but if the reader takes a closer look one finds the scales to be different.


Each line represents one of the 100 trials for the simulations completed for each alternative. The two horizontal lines located in the graphs represent the threshold for gather (upper range of AML) and the target population size (low range of AML). The Most Typical Trial graph includes a dark heavy line (red) which represents what the model has chosen as the trial with the most typical results. This trial closely matches the average of all 100 trials. The most typical trial is useful for making comparisons between alternatives, and for predicting what would be the probable results of the action.

Alternative A: Gather to Low End of AML and Fertility Control Treatment




The results of the modeling for Alternative A indicate that following a 2020 (program default) gather with fertility control that the average population between 2020 and 2030 on a four year cycle would begin to be near AML in 2021 and be within AML by 2024 (gather in 2024) following the 2020 subsequent gather operation and then would stay within AML until approximately 2029. This could potentially make for adjustments in future gather operations and fertility control treatments. This model illustrates that implementing gather and removal simultaneously with fertility control, that the PEDHMA could reach low end of AML within four years, and maintain the population within AML for up to 5 years following the 2024 gather operation.

Alternative B: Gather to Low End of AML and Do Not Use Fertility Control Treatments



The results of the modeling under Alternative B indicate that when the low end of AML (135) remains in the PEDHMA following the initial gather, that the average population would not go over the upper end of the AML until around the year 2023 (four year cycle) but under the modeling another gather would have to take place in order to bring the PEDHMA back down to the low end of AML. The model indicates that AML can be achieved without fertility control, however reaching the low end of the AML is necessary during each gather (completed every four years) without the use of fertility control and the number of wild horses that would need to be gathered and removed at each gather operation would be higher than for Alternative A.

Alternative C: No Action Alternative



Population modeling was completed for the No Action Alternative. The most typical trial was utilized to demonstrate the projected population over time if a gather does not take place. The graph of most typical trial for the gather area is displayed below as a comparison only. The graph clearly shows the continued increase in population size if a gather is not completed.

F.2.2. Growth Rates

Through the model, average population growth rates were obtained for Alternative A (Gather to Low End of AML and Fertility Control Treatment) and Alternative B (Gather to Low End of AML and Do Not Use Fertility Control Treatments) and Alternative C (No Action) out of 100 trials. Growth rates are displayed for the lowest, median and highest trial.

Table 4. HMA - Percent Average Growth Rates in 11 years

Trial	Alternative A: Gather + Fertility Control	Alternative B: Gather Only	Alternative C: No Gather
Lowest	9.2	13.0	14.6
Median	14.6	19.3	19.3
Highest	21.9	25.9	24.2

Population modeling data reflects that the implementation of fertility control treatments could result in reduced growth rate of the wild horse population within the PEDHMA. Growth rate analyzed for the fertility control treatment alternative were lower than when fertility control treatment was not implemented. The model also indicates that growth rates would not be so low as to cause risk to the population should fertility control treatments be implemented.

F.2.3. Population Modeling Summary

To summarize the results obtained by simulating the range of alternatives for the PEDHMA wild horse gather, the following questions can be addressed.

- *Do any of the Alternatives “crash” the population?*
None of the alternatives indicate that a crash is likely to occur to the population. Minimum population levels and growth rates are all within reasonable levels, and adverse impacts to the population are not likely.
- *What effect does fertility control have on population growth rate?*
As expected, the alternative implementing fertility control (Alternative A) reflects the lowest overall growth rates. The growth rates for the PEDHMA proposed for fertility control are lower than the non-fertility control growth rates.
- *What effect do the different alternatives have on the average population size?*
Based on the average median population trial obtained through the population model for the No Action Alternative the herd size would be 2,635 (Alternative C), for the removal only alternative the herd size would be 284 (Alternative B), and for the removal with fertility control treatment alternative the herd size would be 284 (Alternative A), with a possibility of reduced frequency of gather operations and number of wild horses to be gathered and removed depending on effectiveness in application and duration of GonaCon (fertility control treatment preference) within the population.

The No Action Alternative is unacceptable, however, was analyzed for comparison with the other alternatives. Without wild horse gather operation the population has the potential to double approximately every four years.

APPENDIX G. SCIENTIFIC LITERATURE REVIEW

This appendix includes scientific literature reviews addressing five topics: effects of gathers, effects of wild horses and burros on rangeland ecosystems, effects of fertility control vaccines and sex ratio manipulations, effects of sterilization, and effects of intrauterine devices (IUDs). This scientific literature review was compiled by the BLM's Wild Horse and Burro Program Research Coordinator (Paul Griffin, Ph.D.) and is considered current as of October 2020.

G.1. Effects of Gathers on Wild Horses and Burros

Gathering any wild animals into pens has the potential to cause impacts to individual animals. There is also the potential for impacts to individual horses and burros during transportation, short-term holding, long-term holding that take place after a gather. However, BLM follows guidelines to minimize those impacts and ensure humane animal care and high standards of welfare. The following literature review summarizes the limited number of scientific papers and government reports that have examined the effects of gathers and holding on wild horses and burros.

Two early papers, by Hansen and Mosley (2000) and Ashley and Holcomb (2001) examined limited effects of gathers, including behavioral effects and effects on foaling rates. Hansen and Mosley (2000) observed BLM gathers in Idaho and Wyoming. They monitored wild horse behaviors before and after a gather event and compared the behavioral and reproductive outcomes for animals that were gathered by helicopter against those outcomes for animals that were not. This comparison led to the conclusion that gather activities used at that time had no effect on observed wild horse foraging or social behaviors, in terms of time spent resting, feeding, vigilant, traveling, or engaged in agonistic encounters (Hansen and Mosley 2000). Similarly, the authors did not find any statistically significant difference in foaling rates in the year after the gather in comparisons between horses that were captured, those that were chased by a helicopter but evaded capture, or those that were not chased by a helicopter. The authors concluded that the gathers had no deleterious effects on behavior or reproduction. Ashley and Holcomb (2001) conducted observations of reproductive rates at Garfield Flat HMA in Nevada, where horses were gathered in 1993 and 1997, and compared those observations at Granite Range HMA in Nevada, where there was no gather. The authors found that the two gathers had a short-term effect on foaling rates; pregnant mares that were gathered had lower foaling rates than pregnant mares that were not gathered. The authors suggested that BLM make changes to the gather methods used at that time, to minimize the length of time that pregnant mares are held prior to their release back to the range. Since the publications by Hansen and Mosley (2000) and by Ashley and Holcomb (2001), BLM did make changes to reduce the stress that gathered animals, including pregnant females, may experience as a result of gather and removal activities; these measures have been formalized as policy in the comprehensive animal welfare program (BLM IM 2015-151).

A thorough review of gather practices and their effects on wild horses and burros can be found in a 2008 report from the Government Accounting Office. The report found that the BLM had controls in place to help ensure the humane treatment of wild horses and burros (GAO 2008). The controls included SOPs for gather operations, inspections, and data collection to monitor animal welfare. These procedures led to humane treatment during gathers, and in short-term and

long-term holding facilities. The report found that cumulative effects associated with the capture and removal of excess wild horses include gather-related mortality averaged only about 0.5% and approximately 0.7% of the captured animals, on average, are humanely euthanized due to pre-existing conditions (such as lameness or club feet) in accordance with BLM policy. Scasta (2019) found the same overall mortality rate (1.2%) for BLM WH&B gathers in 2010-2019, with a mortality rate of 0.25% caused directly by the gather, and a mortality rate of 0.94% attributable to euthanasia of animals with pre-existing conditions such as blindness or club-footedness. Scasta (2019) summarized mortality rates from 70 BLM WH&B gathers across nine states, from 2010-2019. Records for 28,821 horses and 2,005 burros came from helicopter and bait/water trapping. For wild burro bait / water trapping, mortality rates were 0.05% due to acute injury caused by the gather process, and death for burros with pre-existing conditions was 0.2% (Scasta 2019). For wild horse bait / water trapping, mortality rates were 0.3% due to acute injury, and the mortality rate due to pre-existing conditions was 1.4% (Scasta 2019). For wild horses gathered with the help of helicopters, mortality rates were only slightly lower than for bait / water trapping, with 0.3% due to acute causes, and 0.8% due to pre-existing conditions (Scasta 2019). Scasta (2019) noted that for other wildlife species capture operations, mortality rates above 2% are considered unacceptable and that, by that measure, BLM WH&B "...welfare is being optimized to a level acceptable across other animal handling disciplines."

The GAO report (2008) noted the precautions that BLM takes before gather operations, including screening potential gather sites for environmental and safety concerns, approving facility plans to ensure that there are no hazards to the animals there, and limiting the speeds that animals travel to trap sites. BLM used SOPs for short-term holding facilities (e.g., corrals) that included procedures to minimize excitement of the animals to prevent injury, separating horses by age, sex, and size, regular observation of the animals, and recording information about the animals in a BLM database. The GAO reported that BLM had regular inspections of short-term holding facilities and the animals held there, ensuring that the corral equipment is up to code and that animals are treated with appropriate veterinary care (including that hooves are trimmed adequately to prevent injury). Mortality was found to be about 5% per year associated with transportation, short term holding, and adoption or sale with limitations. The GAO noted that BLM also had controls in place to ensure humane care at long-term holding facilities (i.e., pastures). BLM staff monitor the number of animals, the pasture conditions, winter feeding, and animal health. Veterinarians from the USDA Animal and Plant Health Inspection Service inspect long-term facilities annually, including a full count of animals, with written reports. Contract veterinarians provide animal care at long-term facilities, when needed. Weekly counts provide an incentive for contractors that operate long-term holding facilities to maintain animal health (GAO 2008). Mortality at long-term holding was found to be about 8% per year, on average (GAO 2008). The mortality rates at short-term and long-term holding facilities are comparable to the natural annual mortality rate on the range of about 16% per year for foals (animals under age 1), about 5-10% per year for horses ages 1-10 years, and about 10-25% for animals aged 10-20 years (Ransom et al. 2016).

In 2010, the American Association of Equine Practitioners (AAEP 2011) was invited by the BLM to visit the BLM operations and facilities, spend time on WH&B gathers and evaluate the management of the wild equids. The AAEP Task Force evaluated horses in the BLM Wild Horse and Burro Program through several visits to wild horse gathers, and short- and long-term holding

facilities. The task force was specifically asked to “review animal care and handling within the Wild Horse and Burro Program, and make whatever recommendations, if any, the Association feels may be indicated, and if possible, issue a public statement regarding the care and welfare of animals under BLM management.” In their report (AAEP 2011), the task force concluded “that the care, handling and management practices utilized by the agency are appropriate for this population of horses and generally support the safety, health status and welfare of the animals.”

In June 2010 BLM invited independent observers organized by American Horse Protection Association (AHPA) to observe BLM gathers and document their findings. AHPA engaged four independent credentialed professionals who are academia-based equine veterinarians or equine specialists. Each observer served on a team of two and was tasked specifically to observe the care and handling of the animals for a 3-4-day period during the gather process and submit their findings to AHPA. An Evaluation Checklist was provided to each of the observers that included four sections: Gather Activities; Horse Handling During Gather; Horse Description; and Temporary Holding Facility. The independent group visited three separate gather operations and found that “BLM and contractors are responsible and concerned about the welfare of the horses before, during and after the gather process” and that “gentle and knowledgeable, used acceptable methods for moving horses... demonstrated the ability to review, assess and adapt procedures to ensure the care and well-being of the animals” (Greene et al. 2013).

BLM commissioned the Natural Resources Council of the National Academies of Sciences (NAS) to conduct an independent, technical evaluation of the science, methodology, and technical decision-making approaches of the BLM Wild Horse and Burro Management Program. Among the conclusions of their 2013 report, NAS (2013) concluded that wild horse populations grow at 15-20 percent a year, and that predation will not typically control population growth rates of free-ranging horses. The report (NAS 2013) also noted that, because there are human-created barriers to dispersal and movement (such as fences and highways) and no substantial predator pressure, maintaining a herd within an AML requires removing animals in roundups, also known as gathers, and may require management actions that limit population growth rates. The report (NAS 2013) examined a number of population growth suppression techniques, including the use of sterilization, fertility control vaccines, and sex ratio manipulation.

The effects of gathers as part of feral horse management have also been documented on National Park Service Lands. Since the 1980s, managers at Theodore Roosevelt National Park have used periodic gathers, removals, and auctions to maintain the feral horse herd size at a carrying capacity level of 50 to 90 horses (Amberg et al. 2014). In practical terms, this carrying capacity is equivalent to an AML. Horse herd sizes at those levels were determined to allow for maintenance of certain sensitive forage plant species. Gathers every 3-5 years did not prevent the herd from self-sustaining. The herd continues to grow, to the point that the NPS now uses gathers and removals along with temporary fertility control methods in its feral horse management (Amberg et al. 2014).

Literature Cited, Effects of Gathers

Amberg, S., K. Kilkus, M. Komp, A. Nadeau, K. Stark, L. Danielson, S. Gardner, E. Iverson, E. Norton, and B. Drazkowski. 2014. Theodore Roosevelt: National Park: Natural resource condition assessment. Natural Resource Report NPS/THRO/NRR—2014/776. National Park Service, Fort Collins, Colorado.

- American Association of Equine Practitioners (AAEP). 2011. Bureau of Land Management; BLM Task Force Report.
- Ashley, M.C., and D.W. Holcomb. 2001. Effect of stress induced by gathers and removals on reproductive success of feral horses. *Wildlife Society Bulletin* 29: 248-254
- Bureau of Land Management (BLM). 2015. Comprehensive animal welfare program for wild horse and burro gathers. Instruction Memorandum (IM) 2015-151.
- Government Accountability Office (GAO). 2008. Bureau of Land Management; Effective Long-Term Options Needed to Manage Unadoptable Wild Horses. Report to the Chairman, Committee on Natural Resources, House of Representatives, GAO-09-77.
- Greene, E.A., C.R. Heleski, S.L. Ralston, and C.L. Stull. 2013. Academic assessment of equine welfare during the gather process of the Bureau of Land Management's wild horse and burro program. *Journal of Equine Veterinary Science* 5: 352-353
- Hansen, K.V., and J.C. Mosley. 2000. Effects of roundups on behavior and reproduction of feral horses. *Journal of Range Management* 53: 479-482
- National Research Council of the National Academies of Sciences (NAS). 2013. Using science to improve the BLM wild horse and burro program: a way forward. National Academies Press. Washington, DC.
- Ransom, J.I., L. Lagos, H. Hrabar, H. Mowrazi, D. Ushkhjargal, and N. Spasskaya. 2016. Wild and feral equid population dynamics. Pages 68-86 in J. I. Ransom and P. Kaczensky, eds., *Wild equids; ecology, management and conservation*. Johns Hopkins University Press, Baltimore, Maryland.
- Scasta, J. D. 2019. Mortality and operational attributes relative to feral horse and burro capture techniques based on publicly available data from 2010-2019. *Journal of Equine Veterinary Science*, 102893.

G.2. Effects of Wild Horses and Burros on Rangeland Ecosystems

The presence of wild horses and wild burros can have substantial effects on rangeland ecosystems, and on the capacity for habitat restoration efforts to achieve landscape conservation and restoration goals.

In the biological sense, all free-roaming horses and burros in North America are feral, meaning that they are descendants of domesticated animals brought to the Americas by European colonists. Horses went extinct in the Americas by the end of the Pleistocene, about 10,000 years ago (Webb 1984; MacFadden 2005). Burros evolved in Eurasia (Geigl et al. 2016). The published literature refers to free-roaming horses and burros as either feral or wild. In the ecological context the terms are interchangeable, but the terms 'wild horse' and 'wild burro' are associated with a specific legal status. The following literature review on the effects of wild

horses and burros on rangeland ecosystems draws on scientific studies of feral horses and burros, some of which also have wild horse or wild burro legal status. The following literature review draws on Parts 1 and 2 of the ‘Science framework for conservation and restoration of the sagebrush biome’ interagency report (Chambers et al. 2017, Crist et al. 2019).

Because of the known damage that overpopulated wild horse and burro herds can cause in rangeland ecosystems, the presence of wild horses and burros is considered a threat to Greater sage-grouse habitat quality, particularly in the bird species’ western range (Beever and Aldridge 2011, USFWS 2013). Wild horse population sizes on federal lands have more than doubled in the five years since the USFWS report (2013) was published (BLM 2018). On lands administered by the BLM, there were an estimated 81,951 BLM-administered wild horses and burros as of March 1, 2018, which does not include foals born in 2018. Lands with wild horses and burros are managed for multiple uses, so it can be difficult to parse out their ecological effects. Despite this, scientific studies designed to separate out those effects, which are summarized below, point to conclusions that landscapes with greater wild horse and burro abundance will tend to have lower resilience to disturbance and lower resistance to invasive plants than similar landscapes with herds at or below target AML levels.

In contrast to managed livestock grazing, neither the seasonal timing nor the intensity of wild horse and burro grazing can be managed, except through efforts to manage their numbers and distribution. Wild horses live on the range year round, they roam freely, and wild horse populations have the potential to grow 15-20% per year (Wolfe 1980; Eberhardt et al. 1982; Garrett et al 1991; Dawson 2005; Roelle et al. 2010; Scorolli et al. 2010). Although this annual growth rate may be lower in some areas where mountain lions can take foals (Turner and Morrison 2001, Turner 2015), horses tend to favor use of more open habitats (Schoenecker 2016) that are dominated by grasses and shrubs and where ambush is less likely. Horses can compete with managed livestock in forage selected (Scasta et al. 2016). For the majority of wild horse herds, there is little overall evidence that population growth is significantly affected by predation. As a result of the potential for wild horse populations to grow rapidly, impacts from wild horses on water, soil, vegetation, and native wildlife resources (Davies and Boyd 2019) can increase exponentially unless there is active management to limit their population sizes.

The USFWS (2008), Beever and Aldridge (2011), and Chambers et al (2017) summarize much of the literature that quantifies direct ecosystem effects of wild horse presence. Beever and Aldridge (2011) present a conceptual model that illustrates the effects of wild horses on sagebrush ecosystems. In the Great Basin, areas without wild horses had greater shrub cover, plant cover, species richness, native plant cover, and overall plant biomass, and less cover percentage of grazing-tolerant, unpalatable, and invasive plant species, including cheatgrass, compared to areas with horses (Smith 1986; Beever et al. 2008; Davies et al. 2014; Zeigenfuss et al. 2014; Boyd et al. 2017). There were also measurable increases in soil penetration resistance and erosion, decreases in ant mound and granivorous small mammal densities, and changes in reptile communities (Beever et al. 2003; Beever and Brussard 2004; Beever and Herrick 2006; Ostermann-Kelm et al. 2009). Intensive grazing by horses and other ungulates can damage biological crusts (Belnap et al. 2001). In contrast to domestic livestock grazing, where post-fire grazing rest and deferment can foster recovery, wild horse grazing occurs year-round. These

effects imply that horse presence can have broad effects on ecosystem function that could influence conservation and restoration actions.

Many studies corroborate the general conclusion that wild horses can lead to biologically significant changes in rangeland ecosystems, particularly when their populations are overabundant relative to water and forage resources, and other wildlife living on the landscape (Eldridge et al. 2020). Horses are primarily grazers (Hanley and Hanley 1982), but shrubs – including sagebrush – can represent a large part of a horse’s diet, at least in summer in the Great Basin (Nordquist 2011). Grazing by wild horses can have severe impacts on water source quality, aquatic ecosystems and riparian communities as well (Beever and Brussard 2000; Barnett 2002; Nordquist 2011; USFWS 2008; Earnst et al. 2012; USFWS 2012, Kaweck et al. 2018), sometimes excluding native ungulates from water sources (Ostermann-Kelm et al. 2008; USFWS 2008; Perry et al. 2015; Hall et al. 2016; Gooch et al. 2017; Hall et al. 2018). Impacts to riparian vegetation per individual wild horse can exceed impacts per individual domestic cow (Kaweck et al. 2018). Bird nest survival may be lower in areas with wild horses (Zalba and Cozzani 2004), and bird populations have recovered substantially after livestock and / or wild horses have been removed (Earnst et al. 2005; Earnst et al. 2012; Batchelor et al. 2015). Wild horses can spread non-native plant species, including cheatgrass, and may limit the effectiveness of habitat restoration projects (Beever et al. 2003; Couvreur et al. 2004; Jessop and Anderson 2007; Loydi and Zalba 2009). Riparian and wildlife habitat improvement projects intended to increase the availability of grasses, forbs, riparian habitats, and water will likely attract and be subject to heavy grazing and trampling by wild horses that live in the vicinity of the project. Even after domestic livestock are removed, continued wild horse grazing can cause ongoing detrimental ecosystem effects (USFWS 2008; Davies et al. 2014) which may require several decades for recovery (e.g., Anderson and Inouye 2001).

Most analyses of wild horse effects have contrasted areas with wild horses to areas without, which is a study design that should control for effects of other grazers, but historical or ongoing effects of livestock grazing may be difficult to separate from horse effects in some cases (Davies et al. 2014). Analyses have generally not included horse density as a continuous covariate; therefore, ecosystem effects have not been quantified as a linear function of increasing wild horse density. One exception is an analysis of satellite imagery confirming that varied levels of feral horse biomass were negatively correlated with average plant biomass growth (Ziegenfuss et al. 2014).

Horses require access to large amounts of water; an individual can drink an average of 7.4 gallons of water per day (Groenendyk et al. 1988). Despite a general preference for habitats near water (e.g., Crane et al. 1997), wild horses will routinely commute long distances (e.g., 10+ miles per day) between water sources and palatable vegetation (Hampson et al. 2010).

Wild burros can also substantially affect riparian habitats (e.g., Tiller 1997), native wildlife (e.g., Seegmiller and Ohmart 1981), and have grazing and trampling impacts that are similar to wild horses (Carothers et al. 1976; Hanley and Brady 1977; Douglas and Hurst 1983). Where wild burros and Greater sage-grouse co-occur, burros’ year-round use of low-elevation habitats may lead to a high degree of overlap between burros and Greater sage-grouse (Beever and Aldridge 2011).

Literature Cited; Impacts to Rangeland Ecosystems

- Anderson, J.E., and R.S. Inouye. 2001. Landscape-scale changes in plant species abundance and biodiversity of a sagebrush steppe over 45 years. *Ecological Monographs* 71:531-556.
- Barnett, J. 2002. Monitoring feral horse and burro impacts on habitat, Sheldon National Wildlife Refuge. Unpublished report, Sheldon NWR, Lakeview, Oregon.
- Batchelor, J.L., W.J. Ripple, T.M. Wilson, and L.E. Painter. 2015. Restoration of riparian areas following the removal of cattle in the northwestern Great Basin. *Environmental Management* 55:930-942.
- Beever, E.A. and C.L. Aldridge. 2011. Influences of free-roaming equids on sagebrush ecosystems, with focus on greater sage-grouse. *Studies in Avian Biology* 38:273-290.
- Beever, E.A. and P.F. Brussard. 2000. Examining ecological consequences of feral horse grazing using exclosures. *Western North American Naturalist* 63:236-254.
- Beever, E.A. and J.E. Herrick. 2006. Effects of feral horses in Great Basin landscapes on soils and ants: direct and indirect mechanisms. *Journal of Arid Environments* 66:96-112.
- Beever, E.A., R.J. Tausch, and P.F. Brussard. 2003. Characterizing grazing disturbance in semiarid ecosystems across broad scales, using diverse indices. *Ecological Applications* 13:119-136.
- Beever, E.A., and P.F. Brussard. 2004. Community- and landscape-level responses of reptiles and small mammals to feral-horse grazing in the Great Basin. *Journal of Arid Environments*, 59:271-297.
- Beever, E.A., R.J. Tausch, and W.E. Thogmartin. 2008. Multi-scale responses of vegetation to removal of horse grazing from Great Basin (USA) mountain ranges. *Plant Ecology* 196:163-184.
- Belnap, J., J.H. Kaltenecker, R. Rosentreter, J. Williams, S. Leonard, and D. Eldridge. 2001. Biological soil crusts: ecology and management. *USDI-BLM Technical Reference* 1730-2, 119 pp.
- BLM. 2018. Herd Area and Herd Management Area Statistics.
<https://www.blm.gov/programs/wild-horse-and-burro/about-the-program/program-data>.
- Boyd, C.S., K.W. Davies, and G.H. Collins. 2017. Impacts of feral horse use on herbaceous riparian vegetation within a sagebrush steppe ecosystem. *Rangeland Ecology and Management* 70:411-417.
- Carothers, S.W., M.E. Stitt, and R.R. Johnson. 1976. Feral asses on public lands: an analysis of biotic impact, legal considerations and management alternatives. *North American Wildlife Conference* 41:396-405.

- Chambers, J.C., et al. 2017. Science Framework for Conservation and Restoration of the Sagebrush Biome: Linking the Department of the Interior Secretarial Order 3336 to Long-Term Strategic Conservation Actions. Part 1. Science Basis and Applications. RMRS-GTR-360. Fort Collins, CO: U.S Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Crist, M., et al. 2019. Science Framework for Conservation and Restoration of the Sagebrush Biome: Linking the Department of the Interior Secretarial Order 3336 to Long-Term Strategic Conservation Actions. Part 2. Management applications. Gen. Tech. Rep. RMRS-GTR-389. Fort Collins, CO: U.S Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Couvreux, M., B. Christian, K. Verheyen and M. Hermy. 2004. Large herbivores as mobile links between isolated nature reserves through adhesive seed dispersal. *Applied Vegetation Science* 7:229-236.
- Crane, K.K., M.A. Smith, and D. Reynolds. 1997. Habitat selection patterns of feral horses in south central Wyoming. *Journal of Range Management* 50:374-380.
- Davies, K.W., G. Collins, and C.S. Boyd. 2014. Effects of free-roaming horses on semi-arid rangeland ecosystems: an example from the sagebrush steppe. *Ecosphere* 5:1-14.
- Davies, K.W. and C.S. Boyd. 2019. Ecological effects of free-roaming horses in North American rangelands. *Bioscience* 69:558-565.
- Dawson, M. 2005. The Population Ecology of Feral Horses in the Australian Alps, Management Summary. Unpublished report. Australian Alps Liaison Committee, Canberra.
- Douglas, C.L. and T.L. Hurst. 1993. Review and annotated bibliography of feral burro literature. CPSU/UNLV 044/02, 132 pp.
- Earnst, S.L., J.A. Ballard, and D.S. Dobkin. 2005. Riparian songbird abundance a decade after cattle removal on Hart Mountain and Sheldon National Wildlife Refuges. USDA Forest Service Gen. Tech. Rep. PSW-GTR-191. 550-558 pp.
- Earnst, S.L., D.S. Dobkin, and J.A. Ballard. 2012. Changes in avian and plant communities of aspen woodlands over 12 years after livestock removal in the northwest Great Basin. *Conservation Biology* 26: 862-872.
- Eberhardt, L.L., A.K. Majorowicz and J.A. Wilcox, 1982. Apparent rates of increase for two feral horse herds. *The Journal of Wildlife Management*, pp.367-374.
- Eldridge, D.J., J. Ding, and S. K. Travers. 2020. Feral horse activity reduces environmental quality in ecosystems globally. *Biological Conservation* 241:108367.
- Garrott, R.A., D.B. Siniff, and L.L. Eberhardt. 1991. Growth Rates of Feral Horse Populations. *Journal of Wildlife Management* 55: 641-48.

- Geigl, E.M., S. Bar-David, A. Beja-Pereira, E. Cothran, E. Giulotto, H. Hrabar, T. Toyunsuren, and M. Pruvost. 2016. Genetics and Paleogenetics of Equids. Pages 87-104 in Ransom, J.I. and P. Kaczensky, eds. *Wild Equids: Ecology, Management, and Conservation*.
- Gooch, A.M., S.L. Petersen, G.H. Collins, T.S. Smith, B.R. McMillan, and D.L. Eggett. 2017. The impacts of feral horses on the use of water by pronghorn in the Great Basin. *Journal of Arid Environments* 168:38-43.
- Groenendyk, P., B. English, and I. Abetz. 1988. External balance of water and electrolytes in the horse. *Equine Veterinary Journal* 20:189-193.
- Hall, L.K., R.T. Larsen, M.D. Westover, C.C. Day, R.N. Knight, and B.R. McMillan. 2016. Influence of exotic horses on the use of water by communities of native wildlife in a semi-arid environment. *Journal of Arid Environments* 127:100-105.
- Hall, L.K., R.T. Larsen, R.N. Knight, and B.R. McMillan. 2018. Feral horses influence both spatial and temporal patterns of water use by native ungulates in a semi-arid environment. *Ecosphere* 9(1):e02096
- Hampson, B.A., M.A. de Laat, P.C. Mills and C.C. Pollitt. 2010. Distances travelled by feral horses in ‘outback’ Australia. *Equine Veterinary Journal* 42(s38):582-586.
- Hanley, T.A. and W.W. Brady. 1977. Feral burro impact on a Sonoran Desert range. *Journal of Range Management* 30:374-377.
- Hanley, T. A., and K. A. Hanley. 1982. Food resource partitioning by sympatric ungulates on Great Basin rangeland. *Journal of Range Management* 35(2):152-158.
- Jessop, B.D. and V.J. Anderson. 2007. Cheatgrass invasion in salt desert shrublands: benefits of postfire reclamation. *Rangeland Ecology & Management* 60:235-243.
- Kaweck, M.M., J.P. Severson, and K.L. Launchbaugh. 2018. Impacts of wild horses, cattle, and wildlife on riparian areas in Idaho. *Rangelands* 40:45-52.
- Loydi, A. and S.M. Zalba. 2009. Feral horses dung piles as potential invasion windows for alien plant species in natural grasslands. *Plant Ecology* 201:471-480.
- MacFadden, B.J. 2005. Fossil horses – evidence of evolution. *Science* 307: 1728-1730.
- National Research Council. 2013. Using science to improve the BLM wild horse and burro program: a way forward. National Academies Press, Washington, D.C.
- Nordquist, M. K. 2011. Stable isotope diet reconstruction of feral horses (*Equus caballus*) on the Sheldon National Wildlife Refuge, Nevada, USA. Thesis, Brigham Young University, Provo, Utah.

- Ostermann-Kelm, S., E.R. Atwill, E.S. Rubin, M.C. Jorgensen, and W.M. Boyce. 2008. Interactions between feral horses and desert bighorn sheep at water. *Journal of Mammalogy* 89:459-466.
- Ostermann-Kelm, S.D., E.A. Atwill, E.S. Rubin, L.E. Hendrickson, and W.M. Boyce. 2009. Impacts of feral horses on a desert environment. *BMC Ecology* 9:1-10.
- Perry, N.D., P. Morey and G.S. Miguel. 2015. Dominance of a Natural Water Source by Feral Horses. *The Southwestern Naturalist* 60:390-393.
- Roelle, J.E., F.J. Singer, L.C. Zeigenfuss, J.I. Ransom, L. Coates-Markle, and K.A. Schoenecker. 2010. Demography of the Pryor Mountain wild horses 1993–2007. US Geological Survey Scientific Investigations Report 2010–5125. 31p.
- Scasta, J.D., J.L. Beck and C.J. Angwin. 2016. Meta-Analysis of Diet Composition and Potential Conflict of Wild Horses with Livestock and Wild Ungulates on Western Rangelands of North America. *Rangeland Ecology & Management*.
- Schoenecker, K.A., S.R.B. King, M.K. Nordquist, D. Nandintseg, and Q. Cao. 2016. Habitat and diet of equids. In: *Wild equids: ecology, management, and conservation*, J. I. Ransom and P. Kaczensky, eds. Johns Hopkins University Press. Baltimore, Maryland.
- Scorolli, A.L. and A.C.L. Cazorla. 2010. Demography of feral horses (*Equus caballus*): a long-term study in Tornquist Park, Argentina. *Wildlife Research* 37:207-214.
- Seegmiller, R.F., and R.D. Ohmart. 1981. Ecological relationships of feral burros and desert bighorn sheep. *Wildlife Monographs* 78:3-58.
- Smith, M.A. 1986. Impacts of feral horse grazing on rangelands: an overview. *Journal of Equine Science* 6:236-238.
- Tiller, B.L. 1997. Feral burro populations: distribution and damage assessment. Pacific Northwest National Laboratory 11879. U.S. Army, Department of Public Works, Fort Irwin, California.
- Turner, J.W. and M.L. Morrison. 2001. Influence of predation by mountain lions on numbers and survivorship of a feral horse population. *The Southwestern Naturalist* 46:183-190.
- Turner, J.W. 2015. Environmental influences on movements and distribution of a wild horse (*Equus caballus*) population in western Nevada, USA: a 25-year study. *Journal of Natural History* 49 (39-40):2437-2464.
- USFWS. 2008. Revised, Final Environmental Assessment for Horse and Burro Management at Sheldon National Wildlife Refuge. April 2008. U.S. Fish and Wildlife Service, Lake County, Oregon.

- USFWS. 2012. Sheldon National Wildlife Refuge Comprehensive Conservation Plan. USFWS, Lakeview, Oregon.
- USFWS. 2013. Greater Sage-grouse conservation objectives: final report. U.S. Fish and Wildlife Service, Denver, Colorado. February 2013.
- Webb, S.D. 1989. Ten million years of mammal extinction in North America. In Martin, P.S. and Klein, R.G. eds., Quaternary extinctions: a prehistoric revolution. University of Arizona Press.
- Wolfe, M.L. 1980. Feral horse demography: a preliminary report. *Journal of Range Management* 33:354-360.
- Zalba, S.M., and N.C. Cozzani. 2004. The impact of feral horses on grassland bird communities in Argentina. *Animal Conservation*, 7:35-44.
- Ziegenfuss, L.C., K.A. Schoenecker, J.I. Ransom, D.A. Ignizio, and T. Mask. 2014. Influence of nonnative and native ungulate biomass and seasonal precipitation on vegetation production in a great basin ecosystem. *Western North American Naturalist* 74:286-298.

G.3. Effects of Fertility Control Vaccines and Sex Ratio Manipulations

Various forms of fertility control can be used in wild horses and wild burros, with the goals of maintaining herds at or near AML, reducing fertility rates, and reducing the frequency of gathers and removals. The WFRHBA of 1971 specifically provides for contraception and sterilization (16 U.S.C. 1333 section 3.b.1). Fertility control measures have been shown to be a cost-effective and humane treatment to slow increases in wild horse populations or, when used in combination with gathers, to reduce horse population size (Bartholow 2004, de Seve and Boyles-Griffin 2013, Fonner and Bohara 2017). Although fertility control treatments may be associated with a number of potential physiological, behavioral, demographic, and genetic effects, those impacts are generally minor and transient, do not prevent overall maintenance of a self-sustaining population, and do not generally outweigh the potential benefits of using contraceptive treatments in situations where it is a management goal to reduce population growth rates (Garrott and Oli 2013).

An extensive body of peer-reviewed scientific literature details the impacts of fertility control methods on wild horses and burros. No finding of excess animals is required for BLM to pursue contraception in wild horses or wild burros, but NEPA analysis has been required. This review focuses on peer-reviewed scientific literature. The summary that follows first examines effects of fertility control vaccine use in mares, then of sex ratio manipulation. This review does not examine effects of spaying and neutering. Cited studies are generally limited to those involving horses and burros, except where including studies on other species helps in making inferences about physiological or behavioral questions not yet addressed in horses or burros specifically. While most studies reviewed here refer to horses, burros are extremely similar in terms of physiology, such that expected effects are comparable, except where differences between the species are noted.

On the whole, the identified impacts are generally transient and affect primarily the individuals treated. Fertility control that affects individual horses and burros does not prevent BLM from ensuring that there will be self-sustaining populations of wild horses and burros in single herd management areas (HMAs), in complexes of HMAs, and at regional scales of multiple HMAs and complexes. Under the WFRHBA of 1971, BLM is charged with maintaining self-reproducing populations of wild horses and burros. The National Academies of Sciences (2013) encouraged BLM to manage wild horses and burros at the spatial scale of “metapopulations” – that is, across multiple HMAs and complexes in a region. In fact, many HMAs have historical and ongoing genetic and demographic connections with other HMAs, and BLM routinely moves animals from one to another to improve local herd traits and maintain high genetic diversity. The NAS report (2013) includes information (pairwise genetic ‘fixation index’ values for sampled WH&B herds) confirming that WH&B in the vast majority of HMAs are genetically similar to animals in multiple other HMAs.

All fertility control methods affect the behavior and physiology of treated animals (NAS 2013), and are associated with potential risks and benefits, including effects of handling, frequency of handling, physiological effects, behavioral effects, and reduced population growth rates (Hampton et al. 2015). Contraception alone does not remove excess horses from an HMA’s population, so one or more gathers are usually needed in order to bring the herd down to a level close to AML. Horses are long-lived, potentially reaching 20 years of age or more in the wild. Except in cases where extremely high fractions of mares are rendered infertile over long time periods of (i.e., 10 or more years), fertility control methods such as immunocontraceptive vaccines and sex ratio manipulation are not very effective at reducing population growth rates to the point where births equal deaths in a herd. However, even more modest fertility control activities can reduce the frequency of horse gather activities, and costs to taxpayers. Bartholow (2007) concluded that the application of 2-year or 3-year contraceptives to wild mares could reduce operational costs in a project area by 12-20%, or up to 30% in carefully planned population management programs. Because applying contraception to horses requires capturing and handling, the risks and costs associated with capture and handling of horses may be comparable to those of gathering for removal, but with expectedly lower adoption and long-term holding costs. Population growth suppression becomes less expensive if fertility control is long-lasting (Hobbs et al. 2000).

In the context of BLM wild horse and burro management, fertility control vaccines and sex ratio manipulation rely on reducing the number of reproducing females. Taking into consideration available literature on the subject, the National Academies of Sciences concluded in their 2013 report that forms of fertility control vaccines were two of the three ‘most promising’ available methods for contraception in wild horses and burros (NAS 2013). That report also noted that sex ratio manipulations where herds have approximately 60% males and 40% females can expect lower annual growth rates, simply as a result of having a lower number of reproducing females.

G.3.1. Fertility Control Vaccines

Fertility control vaccines (also known as immunocontraceptives) meet BLM requirements for safety to mares and the environment (EPA 2009a, 2012). Because they work by causing an immune response in treated animals, there is no risk of hormones or toxins being taken into the food chain when a treated mare dies. The BLM and other land managers have mainly used three

fertility control vaccine formulations for fertility control of wild horse mares on the range: ZonaStat-H, PZP-22, and GonaCon-Equine. As other formulations become available, they may be applied in the future.

In any vaccine, the antigen is the stimulant to which the body responds by making antigen-specific antibodies. Those antibodies then signal to the body that a foreign molecule is present, initiating an immune response that removes the molecule or cell. Adjuvants are additional substances that are included in vaccines to elevate the level of immune response. Adjuvants help to incite recruitment of lymphocytes and other immune cells which foster a long-lasting immune response that is specific to the antigen.

Liquid emulsion vaccines can be injected by hand or remotely administered in the field using a pneumatic dart (Roelle and Ransom 2009, Rutberg et al. 2017, McCann et al. 2017) in cases where mares are relatively approachable. Use of remotely delivered (dart-delivered) vaccine is generally limited to populations where individual animals can be accurately identified and repeatedly approached within 50 m (BLM 2010). Booster doses can be safely administered by hand or by dart. Even with repeated booster treatments of the vaccines, it is expected that most mares would eventually return to fertility, though some individual mares treated repeatedly may remain infertile. Once the herd size in a project area is at AML and population growth seems to be stabilized, BLM can make adaptive determinations as to the required frequency of new and booster treatments.

BLM has followed SOPs for fertility control vaccine application (BLM IM 2009-090). Herds selected for fertility control vaccine use should have annual growth rates over 5%, have a herd size over 50 animals, and have a target rate of treatment of between 50% and 90% of female wild horses or burros. The IM requires that treated mares be identifiable via a visible freeze brand or individual color markings, so that their vaccination history can be known. The IM calls for follow-up population surveys to determine the realized annual growth rate in herds treated with fertility control vaccines.

Vaccine Formulations: Porcine Zona Pellucida (PZP)

PZP vaccines have been used on dozens of horse herds by the National Park Service, US Forest Service, Bureau of Land Management, and Native American tribes and PZP vaccine use is approved for free-ranging wild and feral horse herds in the United States (EPA 2012). PZP use can reduce or eliminate the need for gathers and removals, if very high fractions of mares are treated over a very long time period (Turner et al. 1997). PZP vaccines have been used extensively in wild horses (NAS 2013), and in feral burros on Caribbean islands (Turner et al. 1996, French et al. 2017). PZP vaccine formulations are produced as ZonaStat-H, an EPA-registered commercial product (EPA 2012, SCC 2015), as PZP-22, which is a formulation of PZP in polymer pellets that can lead to a longer immune response (Turner et al. 2002, Rutberg et al. 2017), and as Spayvac, where the PZP protein is enveloped in liposomes (Killian et al. 2008, Roelle et al. 2017, Bechert and Fraker 2018). ‘Native’ PZP proteins can be purified from pig ovaries (Liu et al. 1989). Recombinant ZP proteins may be produced with molecular techniques (Gupta and Minhas 2017, Joonè et al. 2017a, Nolan et al. 2018a).

When advisories on the product label (EPA 2015) are followed, the product is safe for users and the environment (EPA 2012). In keeping with the EPA registration for ZonaStat-H (EPA 2012;

reg. no. 86833-1), certification through the Science and Conservation Center in Billings Montana is required to apply that vaccine to equids.

For maximum effectiveness, PZP is administered within the December to February timeframe. When applying ZonaStat-H, first the primer with modified Freund's Complete adjuvant is given and then the booster with Freund's Incomplete adjuvant is given 2-6 weeks later. Preferably, the timing of the booster dose is at least 1-2 weeks prior to the onset of breeding activity. Following the initial 2 inoculations, only annual boosters are required. For the PZP-22 formulation, each released mare would receive a single dose of the two-year PZP contraceptive vaccine at the same time as a dose of the liquid PZP vaccine with modified Freund's Complete adjuvant. The pellets are applied to the mare with a large gauge needle and jab-stick into the hip. Although PZP-22 pellets have been delivered via darting in trial studies (Rutberg et al 2017, Carey et al. 2019), BLM does not plan to use darting for PZP-22 delivery until there is more demonstration that PZP-22 can be reliably delivered via dart.

Vaccine Formulations: Gonadotropin Releasing Hormone (GnRH)

GonaCon (which is produced under the trade name GonaCon-Equine for use in feral horses and burros) is approved for use by authorized federal, state, tribal, public and private personnel, for application to free-ranging wild horse and burro herds in the United States (EPA 2013, 2015). GonaCon has been used on feral horses in Theodore Roosevelt National Park and on wild horses administered by BLM (BLM 2015). GonaCon has been produced by USDA-APHIS (Fort Collins, Colorado) in several different formulations, the history of which is reviewed by Miller et al. (2013). GonaCon vaccines present the recipient with hundreds of copies of GnRH as peptides on the surface of a linked protein that is naturally antigenic because it comes from invertebrate hemocyanin (Miller et al 2013). Early GonaCon formulations linked many copies of GnRH to a protein from the keyhole limpet (GonaCon-KHL), but more recently produced formulations where the GnRH antigen is linked to a protein from the blue mussel (GonaCon-B) proved less expensive and more effective (Miller et al. 2008). GonaCon-Equine is in the category of GonaCon-B vaccines.

As with other contraceptives applied to wild horses, the long-term goal of GonaCon-Equine use is to reduce or eliminate the need for gathers and removals (NAS 2013). GonaCon-Equine contraceptive vaccine is an EPA-approved pesticide (EPA, 2009a) that is relatively inexpensive, meets BLM requirements for safety to mares and the environment, and is produced in a USDA-APHIS laboratory. GonaCon is a pharmaceutical-grade vaccine, including aseptic manufacturing technique to deliver a sterile vaccine product (Miller et al. 2013). If stored at 4° C, the shelf life is 6 months (Miller et al 2013).

Miller et al. (2013) reviewed the vaccine environmental safety and toxicity. When advisories on the product label (EPA 2015) are followed, the product is safe for users and the environment (EPA 2009b). EPA waived a number of tests prior to registering the vaccine, because GonaCon was deemed to pose low risks to the environment, so long as the product label is followed (Wang-Cahill et al., *in press*).

GonaCon-Equine can safely be reapplied as necessary to control the population growth rate; booster dose effects may lead to increased effectiveness of contraception, which is generally the intent. Even after booster treatment of GonaCon-Equine, it is expected that most, if not all, mares would return to fertility at some point. Although the exact timing for the return to fertility in mares boosted more than once with GonaCon-Equine has not been quantified, a prolonged return to fertility would be consistent with the desired effect of using GonaCon (e.g., effective contraception).

The adjuvant used in GonaCon, Adjuvac, generally leads to a milder reaction than Freund's Complete Adjuvant (Powers et al. 2011). Adjuvac contains a small number of killed *Mycobacterium avium* cells (Miller et al. 2008, Miller et al. 2013). The antigen and adjuvant are emulsified in mineral oil, such that they are not all presented to the immune system right after injection. It is thought that the mineral oil emulsion leads to a 'depot effect' that is associated with slow or sustained release of the antigen, and a resulting longer-lasting immune response (Miller et al. 2013). Miller et al. (2008, 2013) have speculated that, in cases where memory-B leukocytes are protected in immune complexes in the lymphatic system, it can lead to years of immune response. Increased doses of vaccine may lead to stronger immune reactions, but only to a certain point; when Yoder and Miller (2010) tested varying doses of GonaCon in prairie dogs, antibody responses to the 200µg and 400µg doses were equal to each other but were both higher than in response to a 100µg dose.

Direct Effects: PZP Vaccines

The historically accepted hypothesis explaining PZP vaccine effectiveness posits that when injected as an antigen in vaccines, PZP causes the mare's immune system to produce antibodies that are specific to zona pellucida proteins on the surface of that mare's eggs. The antibodies bind to the mare's eggs surface proteins (Liu et al. 1989), and effectively block sperm binding and fertilization (Zoo Montana, 2000). Because treated mares do not become pregnant but other ovarian functions remain generally unchanged, PZP can cause a mare to continue having regular estrus cycles throughout the breeding season. More recent observations support a complementary hypothesis, which posits that PZP vaccination causes reductions in ovary size and function (Mask et al. 2015, Joonè et al. 2017b, Joonè et al. 2017c, Nolan et al. 2018b, 2018c). PZP vaccines do not appear to interact with other organ systems, as antibodies specific to PZP protein do not cross-react with tissues outside of the reproductive system (Barber and Fayrer-Hosken 2000).

Research has demonstrated that contraceptive efficacy of an injected liquid PZP vaccine, such as ZonaStat-H, is approximately 90% or more for mares treated twice in the first year (Turner and Kirkpatrick 2002, Turner et al. 2008). The highest success for fertility control has been reported when the vaccine has been applied November through February. High contraceptive rates of 90% or more can be maintained in horses that are given a booster dose annually (Kirkpatrick et al. 1992). Approximately 60% to 85% of mares are successfully contracepted for one year when treated simultaneously with a liquid primer and PZP-22 pellets (Rutberg et al. 2017, Carey et al. 2019). Application of PZP for fertility control would reduce fertility in a large percentage of mares for at least one year (Ransom et al. 2011). The contraceptive result for a single application of the liquid PZP vaccine primer dose along with PZP vaccine pellets (PZP-22), based on winter applications, can be expected to fall in the approximate efficacy ranges as follows (based on

figure 2 in Rutberg et al. 2017). Below, the approximate efficacy is measured as the relative decrease in foaling rate for treated mares, compared to control mares:

Year 1	Year 2	Year 3
0 (developing fetuses come to term)	~30-75%	~20-50%

If mares that have been treated with PZP-22 vaccine pellets subsequently receive a booster dose of either the liquid PZP vaccine or the PZP-22 vaccine pellets, the subsequent contraceptive effect is apparently more pronounced and long-lasting. The approximate efficacy following a booster dose can be expected to be in the following ranges (based on figure 3 in Rutberg et al. 2017).

Year 1	Year 2	Year 3	Year 4
0 (developing fetuses come to term)	~50-90%	~55-75%	~40-75%

The fraction of mares treated in a herd can have a large effect on the realized change in growth rate due to PZP contraception, with an extremely high portion of mares required over many years to be treated to totally prevent population-level growth (e.g., Turner and Kirkpatrick 2002). Gather efficiency does not usually exceed 85% via helicopter, and may be less with bait and water trapping, so there will almost always be a portion of the female population uncaptured that is not treated in any given year. Additionally, some mares may not respond to the fertility control vaccine, but instead will continue to foal normally.

Direct Effects: GnRH Vaccines

GonaCon-Equine is one of several vaccines that have been engineered to create an immune response to the gonadotropin releasing hormone peptide (GnRH). GnRH is a small peptide that plays an important role in signaling the production of other hormones involved in reproduction in both sexes. When combined with an adjuvant, a GnRH vaccine stimulates a persistent immune response resulting in prolonged antibody production against GnRH, the carrier protein, and the adjuvant (Miller et al., 2008). The most direct result of successful GnRH vaccination is that it has the effect of decreasing the level of GnRH signaling in the body, as evidenced by a drop in luteinizing hormone levels, and a cessation of ovulation.

GnRH is highly conserved across mammalian taxa, so some inferences about the mechanism and effects of GonaCon-Equine in horses can be made from studies that used different anti-GnRH vaccines, in horses and other taxa. Other commercially available anti-GnRH vaccines include: Improvac (Imboden et al. 2006, Botha et al. 2008, Janett et al. 2009a, Janett et al. 2009b, Schulman et al. 2013, Dalmau et al. 2015, Nolan et al. 2018c), made in South Africa; Equity (Elhay et al. 2007), made in Australia; Improvest, for use in swine (Bohrer et al. 2014); Repro-BLOC (Boedeker et al. 2011); and Bopriva, for use in cows (Balet et al. 2014). Of these, GonaCon-Equine, Improvac, and Equity are specifically intended for horses. Other anti-GnRH vaccine formulations have also been tested, but did not become trademarked products (e.g.,

Goodloe 1991, Dalin et al 2002, Stout et al. 2003, Donovan et al. 2013, Schaut et al. 2018, Yao et al. 2018). The effectiveness and side-effects of these various anti-GnRH vaccines may not be the same as would be expected from GonaCon-Equine use in horses. Results could differ as a result of differences in the preparation of the GnRH antigen, and the choice of adjuvant used to stimulate the immune response. For some formulations of anti-GnRH vaccines, a booster dose is required to elicit a contraceptive response, though GonaCon can cause short-term contraception in a fraction of treated animals from one dose (Powers et al. 2011, Gionfriddo et al. 2011a, Baker et al. 2013, Miller et al 2013).

GonaCon can provide multiple years of infertility in several wild ungulate species, including horses (Killian et al., 2008; Gray et al., 2010). The lack of estrus cycling that results from successful GonaCon vaccination has been compared to typical winter period of anoestrus in open mares. As anti-GnRH antibodies decline over time, concentrations of available endogenous GnRH increase and treated animals usually regain fertility (Power et al., 2011).

Females that are successfully contracepted by GnRH vaccination enter a state similar to anestrus, have a lack of or incomplete follicle maturation, and no ovarian cycling (Botha et al. 2008, Nolan et al. 2018c). A leading hypothesis is that anti-GnRH antibodies bind GnRH in the hypothalamus – pituitary ‘portal vessels,’ preventing GnRH from binding to GnRH-specific binding sites on gonadotroph cells in the pituitary, thereby limiting the production of gonadotropin hormones, particularly luteinizing hormone (LH) and, to a lesser degree, follicle-stimulating hormone (FSH) (Powers et al. 2011, NAS 2013). This reduction in LH (and FSH), and a corresponding lack of ovulation, has been measured in response to treatment with anti-GnRH vaccines (Boedeker et al. 2011, Garza et al. 1986).

Females successfully treated with anti-GnRH vaccines have reduced progesterone levels (Garza et al. 1986, Stout et al. 2003, Imboden et al. 2006, Elhay 2007, Botha et al. 2008, Killian et al. 2008, Miller et al. 2008, Janett et al. 2009, Schulman et al. 2013, Balet et al 2014, Dalmau et al. 2015) and β -17 estradiol levels (Elhay et al. 2007), but no great decrease in estrogen levels (Balet et al. 2014). Reductions in progesterone do not occur immediately after the primer dose, but can take several weeks or months to develop (Elhay et al. 2007, Botha et al. 2008, Schulman et al. 2013, Dalmau et al. 2015). This indicates that ovulation is not occurring and corpora lutea, formed from post-ovulation follicular tissue, are not being established.

Antibody titer measurements are proximate measures of the antibody concentration in the blood specific to a given antigen. Anti-GnRH titers generally correlate with a suppressed reproduction system (Gionfriddo et al. 2011a, Powers et al. 2011). Various studies have attempted to identify a relationship between anti-GnRH titer levels and infertility, but that relationship has not been universally predictable or consistent. The time length that titer levels stay high appears to correlate with the length of suppressed reproduction (Dalin et al. 2002, Levy et al. 2011, Donovan et al. 2013, Powers et al. 2011). For example, Goodloe (1991) noted that mares did produce elevated titers and had suppressed follicular development for 11-13 weeks after treatment, but that all treated mares ovulated after the titer levels declined. Similarly, Elhay (2007) found that high initial titers correlated with longer-lasting ovarian and behavioral anoestrus. However, Powers et al. (2011) did not identify a threshold level of titer that was consistently indicative of suppressed reproduction despite seeing a strong correlation between

antibody concentration and infertility, nor did Schulman et al. (2013) find a clear relationship between titer levels and mare acyclicity.

In many cases, young animals appear to have higher immune responses, and stronger contraceptive effects of anti-GnRH vaccines than older animals (Brown et al. 1994, Curtis et al. 2001, Stout et al. 2003, Schulman et al. 2013). Vaccinating with GonaCon at too young an age, though, may prevent effectiveness; Gionfriddo et al. (2011a) observed weak effects in 3-4-month-old fawns. It has not been possible to predict which individuals of a given age class will have long-lasting immune responses to the GonaCon vaccine. Gray (2010) noted that mares in poor body condition tended to have lower contraceptive efficacy in response to GonaCon-B. Miller et al. (2013) suggested that higher parasite loads might have explained a lower immune response in free-roaming horses than had been observed in a captive trial. At this time, it is unclear what the most important factors affecting efficacy are.

Several studies have monitored animal health after immunization against GnRH. GonaCon treated mares did not have any measurable difference in uterine edema (Killian 2006, 2008). Powers et al. (2011, 2013) noted no differences in blood chemistry except a mildly elevated fibrinogen level in some GonaCon treated elk. In that study, one sham-treated elk and one GonaCon treated elk each developed leukocytosis, suggesting that there may have been a causal link between the adjuvant and the effect. Curtis et al. (2008) found persistent granulomas at GonaCon-KHL injection sites three years after injection, and reduced ovary weights in treated females. Yoder and Miller (2010) found no difference in blood chemistry between GonaCon treated and control prairie dogs. One of 15 GonaCon treated cats died without explanation, and with no determination about cause of death possible based on necropsy or histology (Levy et al. 2011). Other anti-GnRH vaccine formulations have led to no detectable adverse effects (in elephants; Boedeker et al. 2011), though Imboden et al. (2006) speculated that young treated animals might conceivably have impaired hypothalamic or pituitary function.

Kirkpatrick et al. (2011) raised concerns that anti-GnRH vaccines could lead to adverse effects in other organ systems outside the reproductive system. GnRH receptors have been identified in tissues outside of the pituitary system, including in the testes and placenta (Khodr and Siler-Khodr 1980), ovary (Hsueh and Erickson 1979), bladder (Coit et al. 2009), heart (Dong et al. 2011), and central nervous system, so it is plausible that reductions in circulating GnRH levels could inhibit physiological processes in those organ systems. Kirkpatrick et al. (2011) noted elevated cardiological risks to human patients taking GnRH agonists (such as leuprolide), but the National Academy of Sciences (2013) concluded that the mechanism and results of GnRH agonists would be expected to be different from that of anti-GnRH antibodies; the former flood GnRH receptors, while the latter deprive receptors of GnRH.

Reversibility and Effects on Ovaries: PZP Vaccines

In most cases, PZP contraception appears to be temporary and reversible, with most treated mares returning to fertility over time (Kirkpatrick and Turner 2002). The ZonaStat-H formulation of the vaccine tends to confer only one year of efficacy per dose. Some studies have found that a PZP vaccine in long-lasting pellets (PZP-22) can confer multiple years of contraception (Turner et al. 2007), particularly when boosted with subsequent PZP vaccination

(Rutberg et al. 2017). Other trial data, though, indicate that the pelleted vaccine may only be effective for one year (J. Turner, University of Toledo, Personal Communication to BLM).

The purpose of applying PZP vaccine treatment is to prevent mares from conceiving foals, but BLM acknowledges that long-term infertility, or permanent sterility, could be a result for some number of individual wild horses receiving PZP vaccinations. The rate of long-term or permanent sterility following vaccinations with PZP is hard to predict for individual horses, but that outcome appears to increase in likelihood as the number of doses increases (Kirkpatrick and Turner 2002). Permanent sterility for mares treated consecutively in each of 5-7 years was observed by Nuñez et al. (2010, 2017). In a graduate thesis, Knight (2014) suggested that repeated treatment with as few as three to four years of PZP treatment may lead to longer-term sterility, and that sterility may result from PZP treatment before puberty. Repeated treatment with PZP led long-term infertility in Przewalski's horses receiving as few as one PZP booster dose (Feh 2012). However, even if some number of mares become sterile as a result of PZP treatment, that potential result would be consistent with the contraceptive purpose that motivates BLM's potential use of the vaccine.

In some number of individual mares, PZP vaccination may cause direct effects on ovaries (Gray and Cameron 2010, Joonè et al. 2017b, Joonè et al. 2017c, Joonè et al. 2017d, Nolan et al. 2018b). Joonè et al. (2017a) noted reversible effects on ovaries in mares treated with one primer dose and booster dose. Joonè et al. (2017c) and Nolan et al. (2018b) documented decreased anti-Mullerian hormone (AMH) levels in mares treated with native or recombinant PZP vaccines; AMH levels are thought to be an indicator of ovarian function. Bechert et al. (2013) found that ovarian function was affected by the SpayVac PZP vaccination, but that there were no effects on other organ systems. Mask et al. (2015) demonstrated that equine antibodies that resulted from SpayVac immunization could bind to oocytes, ZP proteins, follicular tissues, and ovarian tissues. It is possible that result is specific to the immune response to SpayVac, which may have lower PZP purity than ZonaStat or PZP-22 (Hall et al. 2016). However, in studies with native ZP proteins and recombinant ZP proteins, Joonè et al. (2017a) found transient effects on ovaries after PZP vaccination in some treated mares; normal estrus cycling had resumed 10 months after the last treatment. SpayVac is a patented formulation of PZP in liposomes that led to multiple years of infertility in some breeding trials (Killian et al. 2008, Roelle et al. 2017, Bechert and Fraker 2018), but unacceptably poor efficacy in a subsequent trial (Kane 2018). Kirkpatrick et al. (1992) noted effects on horse ovaries after three years of treatment with PZP. Observations at Assateague Island National Seashore indicated that the more times a mare is consecutively treated, the longer the time lag before fertility returns, but that even mares treated seven consecutive years did eventually return to ovulation (Kirkpatrick and Turner 2002). Other studies have reported that continued PZP vaccine applications may result in decreased estrogen levels (Kirkpatrick et al. 1992) but that decrease was not biologically significant, as ovulation remained similar between treated and untreated mares (Powell and Monfort 2001). Bagavant et al. (2003) demonstrated T-cell clusters on ovaries, but no loss of ovarian function after ZP protein immunization in macaques.

Reversibility and Effects on Ovaries: GnRH Vaccines

The NAS (2013) review pointed out that single doses of GonaCon-Equine do not lead to high rates of initial effectiveness, or long duration. Initial effectiveness of one dose of GonaCon-

Equine vaccine appears to be lower than for a combined primer plus booster dose of the PZP vaccine Zonastat-H (Kirkpatrick et al. 2011), and the initial effect of a single GonaCon dose can be limited to as little as one breeding season. However, preliminary results on the effects of boosted doses of GonaCon-Equine indicate that it can have high efficacy and longer-lasting effects in free-roaming horses (Baker et al. 2017, 2018) than the one-year effect that is generally expected from a single booster of Zonastat-H.

Too few studies have reported on the various formulations of anti-GnRH vaccines to make generalizations about differences between products, but GonaCon formulations were consistently good at causing loss of fertility in a statistically significant fraction of treated mares for at least one year (Killian et al. 2009, Gray et al. 2010, Baker et al. 2013, 2017, 2018). With few exceptions (e.g., Goodloe 1991), anti-GnRH treated mares gave birth to fewer foals in the first season when there would be an expected contraceptive effect (Botha et al. 2008, Killian et al. 2009, Gray et al. 2010, Baker et al. 2013, 2018). Goodloe (1991) used an anti-GnRH-KHL vaccine with a triple adjuvant, in some cases attempting to deliver the vaccine to horses with a hollow-tipped 'biobullet,' but concluded that the vaccine was not an effective immunocontraceptive in that study.

Not all mares should be expected to respond to the GonaCon-equine vaccine; some number should be expected to continue to become pregnant and give birth to foals. In studies where mares were exposed to stallions, the fraction of treated mares that are effectively contracepted in the year after anti-GnRH vaccination varied from study to study, ranging from ~50% (Baker et al. 2017), to 61% (Gray et al. 2010), to ~90% (Killian et al. 2006, 2008, 2009). Miller et al. (2013) noted lower effectiveness in free-ranging mares (Gray et al. 2010) than captive mares (Killian et al. 2009). Some of these rates are lower than the high rate of effectiveness typically reported for the first year after PZP vaccine treatment (Kirkpatrick et al. 2011). In the one study that tested for a difference, darts and hand-injected GonaCon doses were equally effective in terms of fertility outcome (McCann et al. 2017).

In studies where mares were not exposed to stallions, the duration of effectiveness also varied. A primer and booster dose of Equity led to anoestrus for at least 3 months (Elhay et al. 2007). A primer and booster dose of Improvac also led to loss of ovarian cycling for all mares in the short term (Imboden et al. 2006, Nolan et al. 2018c). It is worth repeating that those vaccines do not have the same formulation as GonaCon.

Results from horses (Baker et al. 2017, 2018) and other species (Curtis et al. 2001) suggest that providing a booster dose of GonaCon-Equine will increase the fraction of temporarily infertile animals to higher levels than would a single vaccine dose alone.

Longer-term infertility has been observed in some mares treated with anti-GnRH vaccines, including GonaCon-Equine. In a single-dose mare captive trial with an initial year effectiveness of 94%, Killian et al. (2008) noted infertility rates of 64%, 57%, and 43% in treated mares during the following three years, while control mares in those years had infertility rates of 25%, 12%, and 0% in those years. GonaCon effectiveness in free-roaming populations was lower, with infertility rates consistently near 60% for three years after a single dose in one study (Gray et al. 2010) and annual infertility rates decreasing over time from 55% to 30% to 0% in another study with one dose (Baker et al. 2017, 2018). Similarly, gradually increasing fertility rates were

observed after single dose treatment with GonaCon in elk (Powers et al. 2011) and deer (Gionfriddo et al. 2011a).

Baker et al. (2017, 2018) observed a return to fertility over 4 years in mares treated once with GonaCon, but then noted extremely low fertility rates of 0% and 16% in the two years after the same mares were given a booster dose four years after the primer dose. Four of nine mares treated with primer and booster doses of Improvac did not return to ovulation within 2 years of the primer dose (Imboden et al. 2006), though one should probably not make conclusions about the long-term effects of GonaCon-Equine based on results from Improvac.

It is difficult to predict which females will exhibit strong or long-term immune responses to anti-GnRH vaccines (Killian et al. 2006, Miller et al. 2008, Levy et al. 2011). A number of factors may influence responses to vaccination, including age, body condition, nutrition, prior immune responses, and genetics (Cooper and Herbert 2001, Curtis et al. 2001, Powers et al. 2011). One apparent trend is that animals that are treated at a younger age, especially before puberty, may have stronger and longer-lasting responses (Brown et al. 1994, Curtis et al. 2001, Stout et al. 2003, Schulman et al. 2013). It is plausible that giving GonaCon-Equine to prepubertal mares will lead to long-lasting infertility, but that has not yet been tested.

To date, short term evaluation of anti-GnRH vaccines, show contraception appears to be temporary and reversible. Killian et al. noted long-term effects of GonaCon in some captive mares (2009). However, Baker et al. (2017) observed horses treated with GonaCon-B return to fertility after they were treated with a single primer dose; after four years, the fertility rate was indistinguishable between treated and control mares. It appears that a single dose of GonaCon results in reversible infertility. If long-term treatment resulted in permanent infertility for some treated mares, such permanent infertility fertility would be consistent with the desired effect of using GonaCon (e.g., effective contraception).

Other anti-GnRH vaccines also have had reversible effects in mares. Elhay (2007) noted a return to ovary functioning over the course of 34 weeks for 10 of 16 mares treated with Equity. That study ended at 34 weeks, so it is not clear when the other six mares would have returned to fertility. Donovan et al. (2013) found that half of mares treated with an anti-GnRH vaccine intended for dogs had returned to fertility after 40 weeks, at which point the study ended. In a study of mares treated with a primer and booster dose of Improvac, 47 of 51 treated mares had returned to ovarian cyclicity within 2 years; younger mares appeared to have longer-lasting effects than older mares (Schulman et al. 2013). Joonè et al. (2017) analyzed samples from the Schulman et al. (2013) study and found no significant decrease in anti-Mullerian hormone (AMH) levels in mares treated with GnRH vaccine. AMH levels are thought to be an indicator of ovarian function, so results from Joonè et al. (2017) support the general view that the anoestrus resulting from GnRH vaccination is physiologically similar to typical winter anoestrus. In a small study with a non-commercial anti-GnRH vaccine (Stout et al. 2003), three of seven treated mares had returned to cyclicity within 8 weeks after delivery of the primer dose, while four others were still suppressed for 12 or more weeks. In elk, Powers et al. (2011) noted that contraception after one dose of GonaCon was reversible. In white-tailed deer, single doses of GonaCon appeared to confer two years of contraception (Miller et al. 2000). Ten of 30 domestic cows treated became pregnant within 30 weeks after the first dose of Bopriva (Balet et al. 2014).

Permanent sterility as a result of single-dose or boosted GonaCon-Equine vaccine, or other anti-GnRH vaccines, has not been recorded, but that may be because no long-term studies have tested for that effect. It is conceivable that some fraction of mares could become sterile after receiving one or more booster doses of GonaCon-Equine. If some fraction of mares treated with GonaCon-Equine were to become sterile, though, that result would be consistent with text of the WFRHBA of 1971, as amended, which allows for sterilization to achieve population goals.

In summary, based on the above results related to fertility effects of GonaCon and other anti-GnRH vaccines, application of a single dose of GonaCon-Equine to gathered or remotely-darted wild horses could be expected to prevent pregnancy in perhaps 30%-60% of mares for one year. Some smaller number of wild mares should be expected to have persistent contraception for a second year, and less still for a third year. Applying one booster dose of GonaCon to previously treated mares may lead to four or more years with relatively high rates (80+%) of additional infertility expected (Baker et al. 2018). There is no data to support speculation regarding efficacy of multiple boosters of GonaCon-Equine; however, given it is formulated as a highly immunogenic long-lasting vaccine, it is reasonable to hypothesize that additional boosters would increase the effectiveness and duration of the vaccine.

GonaCon-Equine only affects the fertility of treated animals; untreated animals will still be expected to give birth. Even under favorable circumstances for population growth suppression, gather efficiency might not exceed 85% via helicopter, and may be less with bait and water trapping. Similarly, not all animals may be approachable for darting. The uncaptured or undarted portion of the female population would still be expected to have normally high fertility rates in any given year, though those rates could go up slightly if contraception in other mares increases forage and water availability.

Changes in hormones associated with anti-GnRH vaccination lead to measurable changes in ovarian structure and function. The volume of ovaries reduced in response to treatment (Garza et al. 1986, Dalin et al. 2002, Imboden et al. 2006, Elhay et al. 2007, Botha et al. 2008, Gionfriddo 2011a, Dalmau et al. 2015). Treatment with an anti-GnRH vaccine changes follicle development (Garza et al. 1986, Stout et al. 2003, Imboden et al. 2006, Elhay et al. 2007, Donovan et al. 2013, Powers et al. 2011, Balet et al. 2014), with the result that ovulation does not occur. A related result is that the ovaries can exhibit less activity and cycle with less regularity or not at all in anti-GnRH vaccine treated females (Goodloe 1991, Dalin et al. 2002, Imboden et al. 2006, Elhay et al. 2007, Janett et al. 2009a, Powers et al. 2011, Donovan et al. 2013). In studies where the vaccine required a booster, hormonal and associated results were generally observed within several weeks after delivery of the booster dose.

Effects on Existing Pregnancies, Foals, and Birth Phenology: PZP Vaccines

Although fetuses are not explicitly protected under the WFRHBA of 1971, as amended, it is prudent to analyze the potential effects of fertility control vaccines on developing fetuses and foals. Any impacts identified in the literature have been found to be transient, and do not influence the future reproductive capacity of offspring born to treated females.

If a mare is already pregnant, the PZP vaccine has not been shown to affect normal development of the fetus or foal, or the hormonal health of the mare with relation to pregnancy (Kirkpatrick and Turner 2003). Studies on Assateague Island (Kirkpatrick and Turner 2002) showed that once

female offspring born to mares treated with PZP during pregnancy eventually breed, they produce healthy, viable foals. It is possible that there may be transitory effects on foals born to mares or jennies treated with PZP. For example, in mice, Sacco et al. (1981) found that antibodies specific to PZP can pass from mother mouse to pup via the placenta or colostrum, but that did not apparently cause any innate immune response in the offspring: the level of those antibodies were undetectable by 116 days after birth. There was no indication in that study that the fertility or ovarian function of those mouse pups was compromised, nor is BLM aware of any such results in horses or burros. Unsubstantiated, speculative connections between PZP treatment and ‘foal stealing’ has not been published in a peer-reviewed study and thus cannot be verified. ‘Foal stealing,’ where a near-term pregnant mare steals a neonate foal from a weaker mare, is unlikely to be a common behavioral result of including spayed mares in a wild horse herd. McDonnell (2012) noted that “foal stealing is rarely observed in horses, except under crowded conditions and synchronization of foaling,” such as in horse feed lots. Those conditions are not likely in the wild, where pregnant mares will be widely distributed across the landscape, and where the expectation is that parturition dates would be distributed across the normal foaling season. Similarly, although Nettles (1997) noted reported stillbirths after PZP treatments in cynomolgus monkeys, those results have not been observed in equids despite extensive use in horses and burros.

On-range observations from 20 years of application to wild horses indicate that PZP application in wild mares does not generally cause mares to give birth to foals out of season or late in the year (Kirkpatrick and Turner 2003). Nuñez’s (2010) research showed that a small number of mares that had previously been treated with PZP foaled later than untreated mares and expressed the concern that this late foaling “may” impact foal survivorship and decrease band stability, or that higher levels of attention from stallions on PZP-treated mares might harm those mares. However, that paper provided no evidence that such impacts on foal survival or mare well-being actually occurred. Rubenstein (1981) called attention to a number of unique ecological features of horse herds on Atlantic barrier islands, such as where Nuñez made observations, which calls into question whether inferences drawn from island herds can be applied to western wild horse herds. Ransom et al. (2013), though, did identify a potential shift in reproductive timing as a possible drawback to prolonged treatment with PZP, stating that treated mares foaled on average 31 days later than non-treated mares. Results from Ransom et al. (2013), however, showed that over 81% of the documented births in that study were between March 1 and June 21, i.e., within the normal, peak, spring foaling season. Ransom et al. (2013) pointedly advised that managers should consider carefully before using fertility control vaccines in small refugia or rare species. Wild horses and burros managed by BLM do not generally occur in isolated refugia, nor are they at all rare species. The US Fish and Wildlife Service denied a petition to list wild horses as endangered (USFWS 2015). Moreover, any effect of shifting birth phenology was not observed uniformly: in two of three PZP-treated wild horse populations studied by Ransom et al. (2013), foaling season of treated mares extended three weeks and 3.5 months, respectively, beyond that of untreated mares. In the other population, the treated mares foaled within the same time period as the untreated mares. Furthermore, Ransom et al. (2013) found no negative impacts on foal survival even with an extended birthing season. If there are shifts in birth phenology, though, it is reasonable to assume that some negative effects on foal survival for a small number of foals might result from particularly severe weather events (Nuñez et al. 2018).

Effects on Existing Pregnancies, Foals, and Birth Phenology: GnRH Vaccines

Although fetuses are not explicitly protected under the WFRHBA of 1971, as amended, it is prudent to analyze the potential effects of fertility control vaccines on developing fetuses and foals. Any impacts identified in the literature have been found to be transient, and do not influence the future reproductive capacity of offspring born to treated females.

GonaCon and other anti-GnRH vaccines can be injected while a female is pregnant (Miller et al. 2000, Powers et al. 2011, Baker et al. 2013) – in such a case, a successfully contracepted mare will be expected to give birth during the following foaling season, but to be infertile during the same year's breeding season. Thus, a mare injected in November of 2018 would not show the contraceptive effect (i.e., no new foal) until spring of 2020.

GonaCon had no apparent effect on pregnancies in progress, foaling success, or the health of offspring, in horses that were immunized in October (Baker et al. 2013), elk immunized 80-100 days into gestation (Powers et al. 2011, 2013), or deer immunized in February (Miller et al. 2000). Kirkpatrick et al. (2011) noted that anti-GnRH immunization is not expected to cause hormonal changes that would lead to abortion in the horse, but this may not be true for the first 6 weeks of pregnancy (NAS 2013). Curtis et al. (2011) noted that GonaCon-KHL treated white tailed deer had lower twinning rates than controls, but speculated that the difference could be due to poorer sperm quality late in the breeding season, when the treated does did become pregnant. Goodloe (1991) found no difference in foal production between treated and control animals.

Offspring of anti-GnRH vaccine treated mothers could exhibit an immune response to GnRH (Khodr and Siler-Khodr 1980), as antibodies from the mother could pass to the offspring through the placenta or colostrum. In the most extensive study of long-term effects of GonaCon immunization on offspring, Powers et al. (2012) monitored 15 elk fawns born to GonaCon treated cows. Of those, 5 had low titers at birth and 10 had high titer levels at birth. All 15 were of normal weight at birth, and developed normal endocrine profiles, hypothalamic GnRH content, pituitary gonadotropin content, gonad structure, and gametogenesis. All the females became pregnant in their second reproductive season, as is typical. All males showed normal development of secondary sexual characteristics. Powers et al. (2012) concluded that suppressing GnRH in the neonatal period did not alter long-term reproductive function in either male or female offspring. Miller et al. (2013) report elevated anti-GnRH antibody titers in fawns born to treated white tailed deer, but those dropped to normal levels in 11 of 12 of those fawns, which came into breeding condition; the remaining fawn was infertile for three years.

Direct effects on foal survival are equivocal in the literature. Goodloe (1991), reported lower foal survival for a small sample of foals born to anti-GnRH treated mares, but she did not assess other possible explanatory factors such as mare social status, age, body condition, or habitat in her analysis (NAS 2013). Gray et al. (2010) found no difference in foal survival in foals born to free-roaming mares treated with GonaCon.

There is little empirical information available to evaluate the effects of GnRH vaccination on foaling phenology, but those effects are likely to be similar to those for PZP vaccine treated mares in which the effects of the vaccine wear off. It is possible that immunocontracepted mares returning to fertility late in the breeding season could give birth to foals at a time that is out of

the normal range (Nuñez et al. 2010, Ransom et al 2013). Curtis et al. (2001) did observe a slightly later fawning date for GonaCon treated deer in the second year after treatment, when some does regained fertility late in the breeding season. In anti-GnRH vaccine trials in free-roaming horses, there were no published differences in mean date of foal production (Goodloe 1991, Gray et al. 2010). Unpublished results from an ongoing study of GonaCon treated free-roaming mares indicate that some degree of seasonal foaling is possible (D. Baker, Colorado State University, personal communication to Paul Griffin, BLM WH&B Research Coordinator). Because of the concern that contraception could lead to shifts in the timing of parturitions for some treated animals, Ransom et al. (2013) advised that managers should consider carefully before using PZP immunocontraception in small refugia or rare species; the same considerations could be advised for use of GonaCon, but wild horses and burros in most areas do not generally occur in isolated refugia, they are not a rare species at the regional, national, or international level, and genetically they represent descendants of domestic livestock with most populations containing few if any unique alleles (NAS 2013). Moreover, in PZP-treated horses that did have some degree of parturition date shift, Ransom et al. (2013) found no negative impacts on foal survival even with an extended birthing season; however, this may be more related to stochastic, inclement weather events than extended foaling seasons. If there were to be a shift in foaling date for some treated mares, the effect on foal survival may depend on severity of weather and local conditions; for example, Ransom et al. (2013) did not find consistent effects across study sites.

Effects of Marking and Injection

Standard practices require that immunocontraceptive-treated animals be readily identifiable, either via brand marks or unique coloration (BLM 2010). Some level of transient stress is likely to result in newly captured mares that do not have markings associated with previous fertility control treatments. It is difficult to compare that level of temporary stress with the long-term stress that can result from food and water limitation on the range (e.g., Creel et al. 2013). Handling may include freeze-marking, for the purpose of identifying that mare and identifying her vaccine treatment history. Under past management practices, captured mares experienced increased stress levels from handling (Ashley and Holcombe 2001), but BLM has instituted guidelines to reduce the sources of handling stress in captured animals (BLM 2015).

Most mares recover from the stress of capture and handling quickly once released back to the range, and none are expected to suffer serious long term effects from the fertility control injections, other than the direct consequence of becoming temporarily infertile. Injection site reactions associated with fertility control treatments are possible in treated mares (Roelle and Ransom 2009, Bechert et al. 2013, French et al. 2017, Baker et al. 2018), but swelling or local reactions at the injection site are expected to be minor in nature. Roelle and Ransom (2009) found that the most time-efficient method for applying PZP is by hand-delivered injection of 2-year pellets when horses are gathered. They observed only two instances of swelling from that technique. Whether injection is by hand or via darting, GonaCon-Equine is associated with some degree of inflammation, swelling, and the potential for abscesses at the injection site (Baker et al. 2013). Swelling or local reactions at the injection site are generally expected to be minor in nature, but some may develop into draining abscesses. Use of remotely delivered vaccine is generally limited to populations where individual animals can be accurately identified and repeatedly approached. The dart-delivered PZP formulation produced injection-site reactions of varying intensity, though none of the observed reactions appeared debilitating to the animals.

(Roelle and Ransom 2009) but that was not observed with dart-delivered GonaCon (McCann et al. 2017). Joonè et al. (2017a) found that injection site reactions had healed in most mares within 3 months after the booster dose, and that they did not affect movement or cause fever.

Long-lasting nodules observed did not appear to change any animal's range of movement or locomotor patterns and in most cases did not appear to differ in magnitude from naturally occurring injuries or scars. Mares treated with one formulation of GnRH-KHL vaccine developed pyogenic abscesses (Goodloe 1991). Miller et al. (2008) noted that the water and oil emulsion in GonaCon will often cause cysts, granulomas, or sterile abscesses at injection sites; in some cases, a sterile abscess may develop into a draining abscess. In elk treated with GonaCon, Powers et al. (2011) noted up to 35% of treated elk had an abscess form, despite the injection sites first being clipped and swabbed with alcohol. Even in studies where swelling and visible abscesses followed GonaCon immunization, the longer-term nodules observed did not appear to change any animal's range of movement or locomotor patterns (Powers et al. 2013, Baker et al. 2017, 2018). The result that other formulations of anti-GnRH vaccine may be associated with less notable injection site reactions in horses may indicate that the adjuvant formulation in GonaCon leads a single dose to cause a stronger immune reaction than the adjuvants used in other anti-GnRH vaccines. Despite that, a booster dose of GonaCon-Equine appears to be more effective than a primer dose alone (Baker et al. 2017). Horses injected in the hip with Improvac showed only transient reactions that disappeared within 6 days in one study (Botha et al. 2008), but stiffness and swelling that lasted 5 days were noted in another study where horses received Improvac in the neck (Imboden et al. 2006). Equity led to transient reactions that resolved within a week in some treated animals (Elhay et al. 2007). Donovan et al. noted no reactions to the canine anti-GnRH vaccine (2013). In cows treated with Bopriva there was a mildly elevated body temperature and mild swelling at injection sites that subsided within 2 weeks (Balet et al. 2014).

Indirect Effects: PZP Vaccines

One expected long-term, indirect effect on wild horses treated with fertility control would be an improvement in their overall health (Turner and Kirkpatrick 2002). Many treated mares would not experience the biological stress of reproduction, foaling and lactation as frequently as untreated mares. The observable measure of improved health is higher body condition scores (Nuñez et al. 2010). After a treated mare returns to fertility, her future foals would be expected to be healthier overall and would benefit from improved nutritional quality in the mare's milk. This is particularly to be expected if there is an improvement in rangeland forage quality at the same time, due to reduced wild horse population size. Past application of fertility control has shown that mares' overall health and body condition remains improved even after fertility resumes. PZP treatment may increase mare survival rates, leading to longer potential lifespan (Turner and Kirkpatrick 2002, Ransom et al. 2014a) that may be as much as 5-10 years (NPS 2008). To the extent that this happens, changes in lifespan and decreased foaling rates could combine to cause changes in overall age structure in a treated herd (i.e., Turner and Kirkpatrick 2002, Roelle et al. 2010), with a greater prevalence of older mares in the herd (Gross 2000, NPS 2008). Observations of mares treated in past gathers showed that many of the treated mares were larger than, maintained higher body condition than, and had larger healthy foals than untreated mares (BLM, anecdotal observations).

Following resumption of fertility, the proportion of mares that conceive and foal could be increased due to their increased fitness; this has been called a ‘rebound effect.’ Elevated fertility rates have been observed after horse gathers and removals (Kirkpatrick and Turner 1991). If repeated contraceptive treatment leads to a prolonged contraceptive effect, then that may minimize or delay the hypothesized rebound effect. Selectively applying contraception to older animals and returning them to the range could reduce long-term holding costs for such horses, which are difficult to adopt, and may reduce the compensatory reproduction that often follows removals (Kirkpatrick and Turner 1991).

Because successful fertility control in a given herd reduces foaling rates and population growth rates, another indirect effect should be to reduce the number of wild horses that have to be removed over time to achieve and maintain the established AML. Contraception may change a herd’s age structure, with a relative increase in the fraction of older animals in the herd (NPS 2008). Reducing the numbers of wild horses that would have to be removed in future gathers could allow for removal of younger, more easily adoptable excess wild horses, and thereby could eliminate the need to send additional excess horses from this area to off-range holding corrals or pastures for long-term holding.

A principle motivation for use of contraceptive vaccines or sex ratio manipulation is to reduce population growth rates and maintain herd sizes at AML. Where successful, this should allow for continued and increased environmental improvements to range conditions within the project area, which would have long-term benefits to wild horse and burro habitat quality, and well-being of animals living on the range. As the population nears or is maintained at the level necessary to achieve a thriving natural ecological balance, vegetation resources would be expected to recover, improving the forage available. With rangeland conditions more closely approaching a thriving natural ecological balance, and with a less concentrated distribution of wild horses and burros, there should also be less trailing and concentrated use of water sources. Lower population density should lead to reduced competition among wild horses using the water sources, and less fighting among horses accessing water sources. Water quality and quantity would continue to improve to the benefit of all rangeland users including wild horses. Wild horses would also have to travel less distance back and forth between water and desirable foraging areas. Among mares in the herd that remain fertile, a higher level of physical health and future reproductive success would be expected in areas where lower horse and burro population sizes lead to increases in water and forage resources. While it is conceivable that widespread and continued treatment with fertility control vaccines could reduce the birth rates of the population to such a point that birth is consistently below mortality, that outcome is not likely unless a very high fraction of the mares present are all treated in almost every year.

Indirect Effects: GnRH Vaccines

As noted above to PZP vaccines, an expected long-term, indirect effect on wild horses treated with fertility control would be an improvement in their overall health. Body condition of anti-GnRH-treated females was equal to or better than that of control females in published studies. Ransom et al. (2014b) observed no difference in mean body condition between GonaCon-B treated mares and controls. Goodloe (1991) found that GnRH-KHL treated mares had higher survival rates than untreated controls. In other species, treated deer had better body condition

than controls (Gionfriddo et al. 2011b), treated cats gained more weight than controls (Levy et al. 2011), as did treated young female pigs (Bohrer et al. 2014).

Following resumption of fertility, the proportion of mares that conceive and foal could be increased due to their increased fitness; this has been called by some a ‘rebound effect.’ Elevated fertility rates have been observed after horse gathers and removals (Kirkpatrick and Turner 1991). If repeated contraceptive treatment leads to a prolonged contraceptive effect, then that may minimize or delay the hypothesized rebound effect. Selectively applying contraception to older animals and returning them to the range could reduce long-term holding costs for such horses, which are difficult to adopt, and could negate the compensatory reproduction that can follow removals (Kirkpatrick and Turner 1991).

Because successful fertility control would reduce foaling rates and population growth rates, another indirect effect would be to reduce the number of wild horses that have to be removed over time to achieve and maintain the established AML. Contraception would be expected to lead to a relative increase in the fraction of older animals in the herd. Reducing the numbers of wild horses that would have to be removed in future gathers could allow for removal of younger, more easily adoptable excess wild horses, and thereby could eliminate the need to send additional excess horses from this area to off-range holding corrals or pastures for long-term holding. Among mares in the herd that remain fertile, a high level of physical health and future reproductive success would be expected because reduced population sizes should lead to more availability of water and forage resources per capita.

Reduced population growth rates and smaller population sizes could also allow for continued and increased environmental improvements to range conditions within the project area, which would have long-term benefits to wild horse habitat quality. As the local horse abundance nears or is maintained at the level necessary to achieve a thriving natural ecological balance, vegetation resources would be expected to recover, improving the forage available to wild horses and wildlife throughout the area. With rangeland conditions more closely approaching a thriving natural ecological balance, and with a less concentrated distribution of wild horses across the range, there should also be less trailing and concentrated use of water sources. Lower population density would be expected to lead to reduced competition among wild horses using the water sources, and less fighting among horses accessing water sources. Water quality and quantity would continue to improve to the benefit of all rangeland users including wild horses. Wild horses would also have to travel less distance back and forth between water and desirable foraging areas. Should GonaCon-Equine treatment, including booster doses, continue into the future, with treatments given on a schedule to maintain a lowered level of fertility in the herd, the chronic cycle of overpopulation and large gathers and removals might no longer occur, but instead a consistent abundance of wild horses could be maintained, resulting in continued improvement of overall habitat conditions and animal health. While it is conceivable that widespread and continued treatment with GonaCon-Equine could reduce the birth rates of the population to such a point that birth is consistently below mortality, that outcome is not likely unless a very high fraction of the mares present are all treated with primer and booster doses, and perhaps repeated booster doses.

Behavioral Effects: PZP Vaccines

Behavioral difference, compared to mares that are fertile, should be considered as potential results of successful contraception. The NAS report (2013) noted that all forms of fertility suppression have effects on mare behavior, mostly because of the lack of pregnancy and foaling, and concluded that fertility control vaccines were among the most promising fertility control methods for wild horses and burros. The resulting impacts may be seen as neutral in the sense that a wide range of natural behaviors is already observable in untreated wild horses, or mildly adverse in the sense that effects are expected to be transient and to not affect all treated animals.

PZP vaccine-treated mares may continue estrus cycles throughout the breeding season. Ransom and Cade (2009) delineated wild horse behaviors. Ransom et al. (2010) found no differences in how PZP-treated and untreated mares allocated their time between feeding, resting, travel, maintenance, and most social behaviors in three populations of wild horses, which is consistent with Powell's (1999) findings in another population. Likewise, body condition of PZP-treated and control mares did not differ between treatment groups in Ransom et al.'s (2010) study. Nuñez (2010) found that PZP-treated mares had higher body condition than control mares in another population, presumably because energy expenditure was reduced by the absence of pregnancy and lactation. Knight (2014) found that PZP-treated mares had better body condition, lived longer and switched harems more frequently, while mares that foaled spent more time concentrating on grazing and lactation and had lower overall body condition.

In two studies involving a total of four wild horse populations, both Nuñez et al. (2009) and Ransom et al. (2010) found that PZP vaccine treated mares were involved in reproductive interactions with stallions more often than control mares, which is not surprising given the evidence that PZP-treated females of other mammal species can regularly demonstrate estrus behavior while contracepted (Shumake and Killian 1997, Heilmann et al. 1998, Curtis et al. 2001, Duncan et al. 2017). There was no evidence, though, that mare welfare was affected by the increased level of herding by stallions noted in Ransom et al. (2010). Nuñez's later analysis (2017) noted no difference in mare reproductive behavior as a function of contraception history.

Ransom et al. (2010) found that control mares were herded by stallions more frequently than PZP-treated mares, and Nuñez et al. (2009, 2014, 2017, 2018) found that PZP-treated mares exhibited higher infidelity to their band stallion during the non-breeding season than control mares. Madosky et al. (2010) and Knight (2014) found this infidelity was also evident during the breeding season in the same population that Nuñez et al. (2009, 2010, 2014, 2017, 2018) studied. Nuñez et al. (2014, 2017, 2018) concluded that PZP-treated mares changing bands more frequently than control mares could lead to band instability. Nuñez et al. (2009), though, cautioned against generalizing from that island population to other herds. Also, despite any potential changes in band infidelity due to PZP vaccination, horses continued to live in social groups with dominant stallions and one or more mares. Nuñez et al. (2014) found elevated levels of fecal cortisol, a marker of physiological stress, in mares that changed bands. The research is inconclusive as to whether all the mares' movements between bands were related to the PZP treatments themselves or the fact that the mares were not nursing a foal, and did not demonstrate any long-term negative consequence of the transiently elevated cortisol levels. Nuñez et al. 2014 wrote that these effects "...may be of limited concern when population reduction is an urgent priority." Nuñez (2018) and Jones et al. (2019, 2020) noted that band stallions of mares that have

received PZP treatment can exhibit changes in behavior and physiology. Nuñez (2018) cautioned that PZP use may limit the ability of mares to return to fertility, but also noted that, “such aggressive treatments may be necessary when rapid reductions in animal numbers are of paramount importance...If the primary management goal is to reduce population size, it is unlikely (and perhaps less important) that managers achieve a balance between population control and the maintenance of more typical feral horse behavior and physiology.”

In contrast to transient stresses, Creel et al. (2013) highlight that variation in population density is one of the most well-established causal factors of chronic activation of the hypothalamic-pituitary-adrenal axis, which mediates stress hormones; high population densities and competition for resources can cause chronic stress. Creel et al. (2013) also state that “...there is little consistent evidence for a negative association between elevated baseline glucocorticoids and fitness.” Band fidelity is not an aspect of wild horse biology that is specifically protected by the WFRHBA of 1971. It is also notable that Ransom et al. (2014b) found higher group fidelity after a herd had been gathered and treated with a contraceptive vaccine; in that case, the researchers postulated that higher fidelity may have been facilitated by the decreased competition for forage after excess horses were removed. At the population level, available research does not provide evidence of the loss of harem structure among any herds treated with PZP. No biologically significant negative impacts on the overall animals or populations overall, long-term welfare or well-being have been established in these studies.

The National Research Council (2013) found that harem changing was not likely to result in serious adverse effects for treated mares:

“The studies on Shackleford Banks (Nuñez et al., 2009; Madosky et al., 2010) suggest that there is an interaction between pregnancy and social cohesion. The importance of harem stability to mare well-being is not clear, but considering the relatively large number of free-ranging mares that have been treated with liquid PZP in a variety of ecological settings, the likelihood of serious adverse effects seem low.”

Nuñez (2010) stated that not all populations will respond similarly to PZP treatment. Differences in habitat, resource availability, and demography among conspecific populations will undoubtedly affect their physiological and behavioral responses to PZP contraception and need to be considered. Kirkpatrick et al. (2010) concluded that: “the larger question is, even if subtle alterations in behavior may occur, this is still far better than the alternative,” and that the “...other victory for horses is that every mare prevented from being removed, by virtue of contraception, is a mare that will only be delaying her reproduction rather than being eliminated permanently from the range. This preserves herd genetics, while gathers and adoption do not.”

The NAS report (2013) provides a comprehensive review of the literature on the behavioral effects of contraception that puts research up to that date by Nuñez et al. (2009, 2010) into the broader context of all of the available scientific literature, and cautions, based on its extensive review of the literature that:

“... in no case can the committee conclude from the published research that the behavior differences observed are due to a particular compound rather than to the fact that treated

animals had no offspring during the study. That must be borne in mind particularly in interpreting long-term impacts of contraception (e.g., repeated years of reproductive “failure” due to contraception).”

Behavioral Effects: GnRH Vaccines

The result that GonaCon treated mares may have suppressed estrous cycles throughout the breeding season can lead treated mares to behave in ways that are functionally similar to pregnant mares. Where it is successful in mares, GonaCon and other anti-GnRH vaccines are expected to induce fewer estrous cycles when compared to non-pregnant control mares. This has been observed in many studies (Garza et al. 1986, Curtis et al. 2001, Dalin et al. 2002, Killian et al. 2006, Dalmau et al. 2015). Females treated with GonaCon had fewer estrous cycles than control or PZP-treated mares (Killian et al. 2006) or deer (Curtis et al. 2001). Thus, any concerns about PZP treated mares receiving more courting and breeding behaviors from stallions (Nuñez et al. 2009, Ransom et al. 2010) are not generally expected to be a concern for mares treated with anti-GnRH vaccines (Botha et al. 2008).

Ransom et al. (2014b) and Baker et al. (2018) found that GonaCon treated mares had similar rates of reproductive behaviors that were similar to those of pregnant mares. Among other potential causes, the reduction in progesterone levels in treated females may lead to a reduction in behaviors associated with reproduction. Despite this, some females treated with GonaCon or other anti-GnRH vaccines did continue to exhibit reproductive behaviors, albeit at irregular intervals and durations (Dalin et al. 2002, Stout et al. 2003, Imboden et al. 2006), which is a result that is similar to spayed (ovariectomized) mares (Asa et al. 1980). Gray et al. (2009a) and Baker et al. (2018) found no difference in sexual behaviors in mares treated with GonaCon and untreated mares. When progesterone levels are low, small changes in estradiol concentration can foster reproductive estrous behaviors (Imboden et al. 2006). Owners of anti-GnRH vaccine treated mares reported a reduced number of estrous-related behaviors under saddle (Donovan et al. 2013). Treated mares may refrain from reproductive behavior even after ovaries return to cyclicity (Elhay et al. 2007). Studies in elk found that GonaCon treated cows had equal levels of precopulatory behaviors as controls (Powers et al. 2011), though bull elk paid more attention to treated cows late in the breeding season, after control cows were already pregnant (Powers et al. 2011).

Stallion herding of mares, and harem switching by mares are two behaviors related to reproduction that might change as a result of contraception. Ransom et al. (2014b) observed a 50% decrease in herding behavior by stallions after the free-roaming horse population at Theodore Roosevelt National Park was reduced via a gather, and mares there were treated with GonaCon-B. The increased harem tending behaviors by stallions were directed to both treated and control mares. It is difficult to separate any effect of GonaCon in this study from changes in horse density and forage following horse removals.

With respect to treatment with GonaCon or other anti-GnRH vaccines, it is probably less likely that treated mares will switch harems at higher rates than untreated animals, because treated mares are similar to pregnant mares in their behaviors (Ransom et al. 2014b). Indeed, Gray et al. (2009a) found no difference in band fidelity in a free-roaming population of horses with GonaCon treated mares, despite differences in foal production between treated and untreated

mares. Ransom et al. (2014b) actually found increased levels of band fidelity after treatment, though this may have been partially a result of changes in overall horse density and forage availability.

Gray et al. (2009) and Ransom et al. (2014b) monitored non-reproductive behaviors in GonaCon treated populations of free-roaming horses. Gray et al. (2009a) found no difference between treated and untreated mares in terms of activity budget, sexual behavior, proximity of mares to stallions, or aggression. Ransom et al. (2014b) found only minimal differences between treated and untreated mare time budgets, but those differences were consistent with differences in the metabolic demands of pregnancy and lactation in untreated mares, as opposed to non-pregnant treated mares.

Genetic Effects of Fertility Control Vaccines

In HMAs where large numbers of wild horses have recent and / or an ongoing influx of breeding animals from other areas with wild or feral horses, contraception is not expected to cause an unacceptable loss of genetic diversity or an unacceptable increase in the inbreeding coefficient. In any diploid population, the loss of genetic diversity through inbreeding or drift can be prevented by large effective breeding population sizes (Wright 1931) or by introducing new potential breeding animals (Mills and Allendorf 1996). The NAS report (2013) recommended that single HMAs should not be considered as isolated genetic populations. Rather, managed herds of wild horses should be considered as components of interacting metapopulations, with the potential for interchange of individuals and genes taking place as a result of both natural and human-facilitated movements. Introducing 1-2 mares every generation (about every 10 years) is a standard management technique that can alleviate potential inbreeding concerns (BLM 2010).

In the last 10 years, there has been a high realized growth rate of wild horses in most areas administered by the BLM, such that most alleles that are present in any given mare are likely to already be well represented in her siblings, cousins, and more distant relatives. With the exception of horses in a small number of well-known HMAs that contain a relatively high fraction of alleles associated with old Spanish horse breeds (NAS 2013), the genetic composition of wild horses in lands administered by the BLM is consistent with admixtures from domestic breeds. As a result, in most HMAs, applying fertility control to a subset of mares is not expected to cause irreparable loss of genetic diversity. Improved longevity and an aging population are expected results of contraceptive treatment that can provide for lengthening generation time; this result would be expected to slow the rate of genetic diversity loss (Hailer et al. 2006). Based on a population model, Gross (2000) found that a strategy to preferentially treat young animals with a contraceptive led to more genetic diversity being retained than either a strategy that preferentially treats older animals, or a strategy with periodic gathers and removals.

Even if it is the case that repeated treatment with a fertility control vaccine may lead to prolonged infertility, or even sterility in some mares, most HMAs have only a low risk of loss of genetic diversity if logistically realistic rates of contraception are applied to mares. Wild horses in most herd management areas are descendants of a diverse range of ancestors coming from many breeds of domestic horses. As such, the existing genetic diversity in the majority of HMAs does not contain unique or historically unusual genetic markers. Past interchange between HMAs, either through natural dispersal or through assisted migration (i.e., human movement of

horses) means that many HMAs are effectively indistinguishable and interchangeable in terms of their genetic composition (i.e., see the table of F_{st} values in NAS 2013). Roelle and Oyler-McCance (2015) used the VORTEX population model to simulate how different rates of mare sterility would influence population persistence and genetic diversity, in populations with high or low starting levels of genetic diversity, various starting population sizes, and various annual population growth rates. Their results show that the risk of the loss of genetic heterozygosity is extremely low except in case where all of the following conditions are met: starting levels of genetic diversity are low, initial population size is 100 or less, the intrinsic population growth rate is low (5% per year), and very large fractions of the female population are permanently sterilized.

It is worth noting that, although maintenance of genetic diversity at the scale of the overall population of wild horses is an intuitive management goal, there are no existing laws or policies that require BLM to maintain genetic diversity at the scale of the individual herd management area or complex. Also, there is no Bureau-wide policy that requires BLM to allow each female in a herd to reproduce before she is treated with contraceptives.

One concern that has been raised with regards to genetic diversity is that treatment with immunocontraceptives could possibly lead to an evolutionary increase in the frequency of individuals whose genetic composition fosters weak immune responses (Cooper and Larson 2006, Ransom et al. 2014a). Many factors influence the strength of a vaccinated individual's immune response, potentially including genetics, but also nutrition, body condition, and prior immune responses to pathogens or other antigens (Powers et al. 2013). This premise is based on an assumption that lack of response to any given fertility control vaccine is a heritable trait, and that the frequency of that trait will increase over time in a population of vaccine-treated animals. Cooper and Herbert (2001) reviewed the topic, in the context of concerns about the long-term effectiveness of immunocontraceptives as a control agent for exotic species in Australia. They argue that immunocontraception could be a strong selective pressure, and that selecting for reproduction in individuals with poor immune response could lead to a general decline in immune function in populations where such evolution takes place. Other authors have also speculated that differences in antibody titer responses could be partially due to genetic differences between animals (Curtis et al. 2001, Herbert and Trigg 2005). However, Magiafolou et al. (2013) clarify that if the variation in immune response is due to environmental factors (i.e., body condition, social rank) and not due to genetic factors, then there will be no expected effect of the immune phenotype on future generations. It is possible that general health, as measured by body condition, can have a causal role in determining immune response, with animals in poor condition demonstrating poor immune reactions (NAS 2013).

Correlations between physical factors and immune response would not preclude, though, that there could also be a heritable response to immunocontraception. In studies not directly related to immunocontraception, immune response has been shown to be heritable (Kean et al. 1994, Sarker et al. 1999). Unfortunately, predictions about the long-term, population-level evolutionary response to immunocontraceptive treatments are speculative at this point, with results likely to depend on several factors, including: the strength of the genetic predisposition to not respond to the fertility control vaccine; the heritability of that gene or genes; the initial prevalence of that gene or genes; the number of mares treated with a primer dose of the vaccine (which generally

has a short-acting effect); the number of mares treated with one or more booster doses of the vaccine; and the actual size of the genetically-interacting metapopulation of horses within which the vaccine treatment takes place.

BLM is not aware of any studies that have quantified the heritability of a lack of response to immunocontraception such as PZP vaccine or GonaCon-Equine in horses or burros. At this point, there are no studies available from which one could make conclusions about the long-term effects of sustained and widespread immunocontraception treatments on population-wide immune function. Although a few, generally isolated, feral horse populations have been treated with high fractions of mares receiving PZP immunocontraception for long-term population control (e.g., Assateague Island National Park, and Pryor Mountains Herd Management Area), no studies have tested for changes in immune competence in those areas. Relative to the large number of free-roaming feral horses in the western United States, immunocontraception has not been, and is not expected to be used in the type of widespread or prolonged manner that might be required to cause a detectable evolutionary response.

Sex Ratio Manipulation

Skewing the sex ratio of a herd so that there are more males than females is an established BLM management technique for reducing population growth rates. As part of a wild horse and burro gather process, the number of animals returned to the range may include more males, the number removed from the range may include more females, or both. By reducing the proportion of breeding females in a population (as a fraction of the total number of animals present), the technique leads to fewer foals being born, relative to the total herd size.

Sex ratio is typically adjusted in such a way that 60 percent of the horses are male. In the absence of other fertility control treatments, this 60:40 sex ratio can temporarily reduce population growth rates from approximately 20% to approximately 15% (Bartholow 2004). While such a decrease in growth rate may not appear to be large or long-lasting, the net result can be that fewer foals being born, at least for a few years – this can extend the time between gathers, and reduce impacts on-range, and costs off-range. Any impacts of sex ratio manipulation are expected to be temporary because the sex ratio of wild horse and burro foals at birth is approximately equal between males and females (NAS 2013), and it is common for female foals to reproduce by their second year (NAS 2013). Thus, within a few years after a gather and selective removal that leads to more males than females, the sex ratio of reproducing wild horses and burros will be returning toward a 50:50 ratio.

Having a larger number of males than females is expected to lead to several demographic and behavioral changes as noted in the NAS report (2013), including the following. Having more fertile males than females should not alter the fecundity of fertile females. Wild mares may be distributed in a larger number of smaller harems. Competition and aggression between males may cause a decline in male body condition. Female foraging may be somewhat disrupted by elevated male-male aggression. With a greater number of males available to choose from, females may have opportunities to select more genetically fit sires. There would also be an increase in the genetic effective population size because more stallions would be breeding and existing females would be distributed among many more small harems. This last beneficial impact is one reason that skewing the sex ratio to favor males is listed in the BLM wild horse

and burro handbook (BLM 2010) as a method to consider in herds where there may be concern about the loss of genetic diversity; having more males fosters a greater retention of genetic diversity.

Infanticide is a natural behavior that has been observed in wild equids (Feh and Munktuya 2008, Gray 2009), but there are no published accounts of infanticide rates increasing as a result of having a skewed sex ratio in wild horse or wild burro herds. Any comment that implies such an impact would be speculative.

The BLM wild horse and burro management handbook (BLM 2010) discusses this method. The handbook acknowledges that there may be some behavioral impacts of having more males than females. The handbook includes guidelines for when the method should be applied, specifying that this method should be considered where the low end of the AML is 150 animals or greater, and with the result that males comprise 60-70 percent of the herd. Having more than 70 percent males may result in unacceptable impacts in terms of elevated male-male aggression. In NEPA analyses, BLM has chosen to follow these guidelines in some cases, for example:

- In the 2015 Cold Springs HMA Population Management Plan EA (DOI-BLM-V040-2015-022), the low end of AML was 75. Under the preferred alternative, 37 mares and 38 stallions would remain on the HMA. This is well below the 150 head threshold noted above.
- In the 2017 Hog Creek HMA Population Management Plan EA (DOI-BLM-ORWA-V000-2017-0026-EA), BLM clearly identified that maintaining a 50:50 sex ratio was appropriate because the herd size at the low end of AML was only 30 animals.

It is relatively straightforward to speed the return of skewed sex ratios back to a 50:50 ratio. The BLM wild horse and burro handbook (BLM 2010) specifies that, if post-treatment monitoring reveals negative impacts to breeding harems due to sex ratio manipulation, then mitigation measures could include removing males, not introducing additional males, or releasing a larger proportion of females during the next gather.

Literature Cited; Fertility Control Vaccines and Sex Ratio Manipulation

- Asa, C.S., D.A. Goldfoot, M.C. Garcia, and O.J. Ginther. 1980. Sexual behavior in ovariectomized and seasonally anovulatory pony mares (*Equus caballus*). *Hormones and Behavior* 14:46-54.
- Ashley, M.C., and D.W. Holcombe. 2001. Effects of stress induced by gathers and removals on reproductive success of feral horses. *Wildlife Society Bulletin* 29:248-254.
- Baker, D.L., J.G. Powers, M.O. Oehler, J.I. Ransom, J. Gionfriddo, and T.M. Nett. 2013. Field evaluation of the Immunocontraceptive GonaCon-B in Free-ranging Horses (*Equus caballus*) at Theodore Roosevelt National Park. *Journal of Zoo and Wildlife Medicine* 44:S141-S153.
- Baker, D.L., J.G. Powers, J. Ransom, B. McCann, M. Oehler, J. Bruemmer, N. Galloway, D. Eckery, and T. Nett. 2017. Gonadotropin-releasing hormone vaccine (GonaCon-Equine)

- suppresses fertility in free-ranging horses (*Equus caballus*): limitations and side effects. Proceedings of the 8th International Wildlife Fertility Control Conference, Washington, D.C.
- Baker D.L., J.G. Powers, J.I. Ransom, B.E. McCann, M.W. Oehler, J.E. Bruemmer, N.L. Galloway, D. C. Eckery, and T. M. Nett. 2018. Reimmunization increases contraceptive effectiveness of gonadotropin-releasing hormone vaccine (GonaCon-Equine) in free-ranging horses (*Equus caballus*): Limitations and side effects..PLoS ONE 13(7): e0201570.
- Balet, L., F. Janett, J. Hüsler, M. Piechotta, R. Howard, S. Amatayakul-Chantler, A. Steiner, and G. Hirsbrunner, 2014. Immunization against gonadotropin-releasing hormone in dairy cattle: Antibody titers, ovarian function, hormonal levels, and reversibility. *Journal of Dairy Science* 97:2193-2203.
- Bagavant, H., C. Sharp, B. Kurth, and K.S.K. Tung. 2002. Induction and immunohistology of autoimmune ovarian disease in cynomolgus macaques (*Macaca fascicularis*). *American Journal of Pathology* 160:141-149.
- Barber, M.R., and R.A. Fayer-Hosken. 2000. Evaluation of somatic and reproductive immunotoxic effects of the porcine zona pellucida vaccination. *Journal of Experimental Zoology* 286:641-646.
- Bartholow, J.M. 2004. An economic analysis of alternative fertility control and associated management techniques for three BLM wild horse herds. USGS Open-File Report 2004-1199.
- Bartholow, J. 2007. Economic benefit of fertility control in wild horse populations. *The Journal of Wildlife Management* 71:2811-2819.
- Bechert, U., J. Bartell, M. Kutzler, A. Menino, R. Bildfell, M. Anderson, and M. Fraker. 2013. Effects of two porcine zona pellucida immunocontraceptive vaccines on ovarian activity in horses. *The Journal of Wildlife Management* 77:1386-1400.
- Bechert, U.S., and M.A. Fraker. 2018. Twenty years of SpayVac research: potential implications for regulating feral horse and burro populations in the United States. *Human-Wildlife Interactions* 12:117-130.
- Boedeker, N.C., L.A.C. Hayek, S. Murray, D.M. De Avila, and J.L. Brown. 2012. Effects of a gonadotropin-releasing hormone vaccine on ovarian cyclicity and uterine morphology of an Asian elephant (*Elephas maximus*). *Journal of Zoo and Wildlife Medicine* 43:603-614.
- Bohrer, B.M., W.L. Flowers, J.M. Kyle, S.S. Johnson, V.L. King, J.L. Spruill, D.P. Thompson, A.L. Schroeder, and D.D. Boler. 2014. Effect of gonadotropin releasing factor suppression with an immunological on growth performance, estrus activity, carcass characteristics, and meat quality of market gilts. *Journal of Animal Science* 92:4719-4724.

- Botha, A.E., M.L. Schulman, H.J. Bertschinger, A.J. Guthrie, C.H. Annandale, and S.B. Hughes. 2008. The use of a GnRH vaccine to suppress mare ovarian activity in a large group of mares under field conditions. *Wildlife Research* 35:548-554.
- Brown, B.W., P.E. Mattner, P.A. Carroll, E.J. Holland, D.R. Paull, R.M. Hoskinson, and R.D.G. Rigby. 1994. Immunization of sheep against GnRH early in life: effects on reproductive function and hormones in rams. *Journal of Reproduction and Fertility* 101:15-21.
- Bureau of Land Management (BLM). 2010. BLM-4700-1 Wild Horses and Burros Management Handbook. Washington, D.C.
- Bureau of Land Management (BLM). 2015. Instruction Memorandum 2015-151; Comprehensive animal welfare program for wild horse and burro gathers. Washington, D.C.
- Carey, K.A., A. Ortiz, K. Grams, D. Elkins, J.W. Turner, and A.T. Rutberg. 2019. *Wildlife Research* 46:713-718.
- Coit, V.A., F.J. Dowell, and N.P. Evans. 2009. Neutering affects mRNA expression levels for the LH- and GnRH-receptors in the canine urinary bladder. *Theriogenology* 71:239-247.
- Curtis, P.D., R.L. Pooler, M.E. Richmond, L.A. Miller, G.F. Mattfeld, and F.W. Quimby. 2008. Physiological Effects of gonadotropin-releasing hormone immunocontraception in white-tailed deer. *Human-Wildlife Conflicts* 2:68-79.
- Cooper, D.W. and C.A. Herbert. 2001. Genetics, biotechnology and population management of over-abundant mammalian wildlife in Australasia. *Reproduction, Fertility and Development*, 13:451-458.
- Cooper, D.W. and E. Larsen. 2006. Immunocontraception of mammalian wildlife: ecological and immunogenetic issues. *Reproduction*, 132, 821–828.
- Creel, S., B. Dantzer, W. Goymann, and D.R. Rubenstein. 2013. The ecology of stress: effects of the social environment. *Functional Ecology* 27:66-80.
- Curtis, P.D., R.L. Pooler, M.E. Richmond, L.A. Miller, G.F. Mattfeld, and F.W. Quimby. 2001. Comparative effects of GnRH and porcine zona pellucida (PZP) immunocontraceptive vaccines for controlling reproduction in white-tailed deer (*Odocoileus virginianus*). *Reproduction (Cambridge, England) Supplement* 60:131-141.
- Dalmau, A., A. Velarde, P. Rodríguez, C. Pedernera, P. Llonch, E. Fàbrega, N. Casal, E. Mainau, M. Gispert, V. King, and N. Sloomans. 2015. Use of an anti-GnRF vaccine to suppress estrus in crossbred Iberian female pigs. *Theriogenology* 84:342-347.
- Dalin, A.M., Ø. Andresen, and L. Malmgren. 2002. Immunization against GnRH in mature mares: antibody titres, ovarian function, hormonal levels and oestrous behaviour. *Journal of Veterinary Medicine Series A* 49:125-131.

- de Seve, C.W. and S.L. Boyles-Griffin. 2013. An economic model demonstrating the long-term cost benefits of incorporating fertility control into wild horse (*Equus caballus*) management in the United States. *Journal of Zoo and Wildlife Medicine* 44(4s:S34-S37).
- Dong, F., D.C. Skinner, T. John Wu, and J. Ren. 2011. The Heart: A Novel Gonadotrophin-Releasing Hormone Target. *Journal of Neuroendocrinology* 23:456-463.
- Donovan, C.E., T. Hazzard, A. Schmidt, J. LeMieux, F. Hathaway, and M.A. Kutzler. 2013. Effects of a commercial canine gonadotropin releasing hormone vaccine on estrus suppression and estrous behavior in mares. *Animal Reproduction Science*, 142:42-47.
- Duncan, C.L., J.L. King, and P. Stapp. 2017. Effects of prolonged immunocontraception on the breeding behavior of American bison. *Journal of Mammalogy* 98:1272-1287.
- Elhay, M., A. Newbold, A. Britton, P. Turley, K. Dowsett, and J. Walker. 2007. Suppression of behavioural and physiological oestrus in the mare by vaccination against GnRH. *Australian Veterinary Journal* 85:39-45.
- Environmental Protection Agency (EPA). 2009a. Pesticide Fact Sheet: Mammalian Gonadotropin Releasing Hormone (GnRH), New Chemical, Nonfood Use, USEPA-OPP, Pesticides and Toxic Substances. US Environmental Protection Agency, Washington, DC
- Environmental Protection Agency (EPA). 2009b. Memorandum on GonaCon™ Immunocontraceptive Vaccine for Use in White-Tailed Deer. Section 3 Registration. US Environmental Protection Agency, Washington, DC.
- Environmental Protection Agency (EPA). 2012. Porcine Zona Pellucida. Pesticide fact Sheet. Office of Chemical Safety and Pollution Prevention 7505P. 9 pages.
- Environmental Protection Agency (EPA). 2013. Notice of pesticide registration for GonaCon-Equine. US Environmental Protection Agency, Washington, DC.
- Environmental Protection Agency (EPA). 2015. Label and CSF Amendment. November 19, 2015 memo and attachment from Marianne Lewis to David Reinhold. US Environmental Protection Agency, Washington, DC.
- Environmental Protection Agency (EPA). 2012. Porcine Zona Pellucida. Pesticide fact Sheet. Office of Chemical Safety and Pollution Prevention 7505P. 9 pages.
- Feh, C. 2012. Delayed reversibility of PZP (porcine zona pellucida) in free-ranging Przewalski's horse mares. In International Wild Equid Conference. Vienna, Austria: University of Veterinary Medicine.
- Feh, C., and B. Munkhtuya. 2008. Male infanticide and paternity analyses in a socially natural herd of Przewalski's horses: Sexual selection? *Behavioral Processes* 78:335-339.

- Fonner, R. and A.K. Bohara. 2017. Optimal control of wild horse populations with nonlethal methods. *Land Economics* 93:390-412.
- French, H., E. Peterson, R. Ambrosia, H. Bertschinger, M. Schulman, M. Crampton, R. Roth, P. Van Zyl, N. Cameron-Blake, M. Vandenplas, and D. Knobel. 2017. Porcine and recombinant zona pellucida vaccines as immunocontraceptives for donkeys in the Caribbean. *Proceedings of the 8th International Wildlife Fertility Control Conference*, Washington, D.C.
- Garrott, R.A., and M.K. Oli. 2013. A Critical Crossroad for BLM's Wild Horse Program. *Science* 341:847-848.
- Garza, F., D.L. Thompson, D.D. French, J.J. Wiest, R.L. St George, K.B. Ashley, L.S. Jones, P.S. Mitchell, and D.R. McNeill. 1986. Active immunization of intact mares against gonadotropin-releasing hormone: differential effects on secretion of luteinizing hormone and follicle-stimulating hormone. *Biology of Reproduction* 35:347-352.
- Gionfriddo, J.P., A.J. Denicola, L.A. Miller, and K.A. Fagerstone. 2011a. Efficacy of GnRH immunocontraception of wild white-tailed deer in New Jersey. *Wildlife Society Bulletin* 35:142-148.
- Gionfriddo, J.P., A.J. Denicola, L.A. Miller, and K.A. Fagerstone. 2011b. Health effects of GnRH immunocontraception of wild white-tailed deer in New Jersey. *Wildlife Society Bulletin* 35:149-160.
- Goodloe, R.B., 1991. Immunocontraception, genetic management, and demography of feral horses on four eastern US barrier islands. UMI Dissertation Services.
- Gray, M.E. 2009a. The influence of reproduction and fertility manipulation on the social behavior of feral horses (*Equus caballus*). Dissertation. University of Nevada, Reno.
- Gray, M.E. 2009b. An infanticide attempt by a free-roaming feral stallion (*Equus caballus*). *Biology Letters* 5:23-25.
- Gray, M.E., D.S. Thain, E.Z. Cameron, and L.A. Miller. 2010. Multi-year fertility reduction in free-roaming feral horses with single-injection immunocontraceptive formulations. *Wildlife Research* 37:475-481.
- Gray, M.E. and E.Z. Cameron. 2010. Does contraceptive treatment in wildlife result in side effects? A review of quantitative and anecdotal evidence. *Reproduction* 139:45-55.
- Gross, J.E. 2000. A dynamic simulation model for evaluating effects of removal and contraception on genetic variation and demography of Pryor Mountain wild horses. *Biological Conservation* 96:319-330.
- Gupta, S., and V. Minhas. 2017. Wildlife population management: are contraceptive vaccines a feasible proposition? *Frontiers in Bioscience, Scholar* 9:357-374.

- Hailer, F., B. Helander, A.O. Folkestad, S.A. Ganusevich, S. Garstad, P. Hauff, C. Koren, T. Nygård, V. Volke, C. Vilà, and H. Ellegren. 2006. Bottlenecked but long-lived: high genetic diversity retained in white-tailed eagles upon recovery from population decline. *Biology Letters* 2:316-319.
- Hall, S. E., B. Nixon, and R.J. Aiken. 2016. Non-surgical sterilization methods may offer a sustainable solution to feral horse (*Equus caballus*) overpopulation. *Reproduction, Fertility and Development*, published online: <https://doi.org/10.1071/RD16200>
- Hampton, J.O., T.H. Hyndman, A. Barnes, and T. Collins. 2015. Is wildlife fertility control always humane? *Animals* 5:1047-1071.
- Heilmann, T.J., R.A. Garrott, L.L. Cadwell, and B.L. Tiller, 1998. Behavioral response of free-ranging elk treated with an immunocontraceptive vaccine. *Journal of Wildlife Management* 62: 243-250.
- Herbert, C.A. and T.E. Trigg. 2005. Applications of GnRH in the control and management of fertility in female animals. *Animal Reproduction Science*, 88:141-153.
- Hobbs, N.T., D.C. Bowden and D.L. Baker. 2000. Effects of Fertility Control on Populations of Ungulates: General, Stage-Structured Models. *Journal of Wildlife Management* 64:473-491.
- Hsueh, A.J.W. and G.F. Erickson. 1979. Extrapituitary action of gonadotropin-releasing hormone: direct inhibition ovarian steroidogenesis. *Science* 204:854-855.
- Imboden, I., F. Janett, D. Burger, M.A. Crowe, M. Hässig, and R. Thun. 2006. Influence of immunization against GnRH on reproductive cyclicity and estrous behavior in the mare. *Theriogenology* 66:1866-1875.
- Janett, F., U. Lanker, H. Jörg, E. Meijerink, and R. Thun. 2009a. Suppression of reproductive cyclicity by active immunization against GnRH in the adult ewe. *Schweizer Archiv für Tierheilkunde* 151:53-59.
- Janett, F., R. Stump, D. Burger, and R. Thun. 2009b. Suppression of testicular function and sexual behavior by vaccination against GnRH (Equity™) in the adult stallion. *Animal Reproduction Science* 115:88-102.
- Jones, M.M., and C.M.V. Nuñez. 2019. Decreased female fidelity alters male behavior in a feral horse population managed with immunocontraception. *Applied Animal Behaviour Science* 214:34-41.
- Jones, M.M., L. Proops, and C.M.V. Nuñez. 2020. Rising up to the challenge of their rivals: mare infidelity intensifies stallion response to playback of aggressive conspecific vocalizations. *Applied Animal Behaviour Science* (in press): 104949.

- Joonè, C.J., H.J. Bertschinger, S.K. Gupta, G.T. Fosgate, A.P. Arukha, V. Minhas, E. Dieterman, and M.L. Schulman. 2017a. Ovarian function and pregnancy outcome in pony mares following immunocontraception with native and recombinant porcine zona pellucida vaccines. *Equine Veterinary Journal* 49:189-195.
- Joonè, C.J., H. French, D. Knobel, H.J. Bertschinger, and M.L. Schulman. 2017b. Ovarian suppression following PZP vaccination in pony mares and donkey jennies. *Proceedings of the 8th International Wildlife Fertility Control Conference*, Washington, D.C.
- Joonè, C.J., M.L. Schulman, G.T. Fosgate, A.N. Claes, S.K. Gupta, A.E. Botha, A-M Human, and H.J. Bertschinger. 2017c. Serum anti-Müllerian hormone dynamics in mares following immunocontraception with anti-zona pellucida or -GnRH vaccines, *Theriogenology* (2017), doi: 10.1016/
- Joonè, C.J., M.L. Schulman, and H.J. Bertschinger. 2017d. Ovarian dysfunction associated with zona pellucida-based immunocontraceptive vaccines. *Theriogenology* 89:329-337.
- Kane, A.J. 2018. A review of contemporary contraceptives and sterilization techniques for feral horses. *Human-Wildlife Interactions* 12:111-116.
- Kaur, K. and V. Prabha. 2014. Immunocontraceptives: new approaches to fertility control. *BioMed Research International* v. 2014, ArticleID 868196, 15 pp.
<http://dx.doi.org/10.1155/2014/868196>
- Kean, R.P., A. Cahaner, A.E. Freeman, and S.J. Lamont. 1994. Direct and correlated responses to multitrait, divergent selection for immunocompetence. *Poultry Science* 73:18-32.
- Killian, G., N.K. Diehl, L. Miller, J. Rhyon, and D. Thain. 2006. Long-term efficacy of three contraceptive approaches for population control of wild horses. In *Proceedings-Vertebrate Pest Conference*.
- Killian, G., D. Thain, N.K. Diehl, J. Rhyon, and L. Miller. 2008. Four-year contraception rates of mares treated with single-injection porcine zona pellucida and GnRH vaccines and intrauterine devices. *Wildlife Research* 35:531-539.
- Killian, G., T.J. Kreeger, J. Rhyon, K. Fagerstone, and L. Miller. 2009. Observations on the use of GonaCon™ in captive female elk (*Cervus elaphus*). *Journal of Wildlife Diseases* 45:184-188.
- Kirkpatrick, J.F. and J.W. Turner. 1991. Compensatory reproduction in feral horses. *Journal of Wildlife Management* 55:649-652.
- Kirkpatrick, J.F., I.M.K. Liu, J.W. Turner, R. Naugle, and R. Keiper. 1992. Long-term effects of porcine zona pellucida immunocontraception on ovarian function in feral horses (*Equus caballus*). *Journal of Reproduction and Fertility* 94:437-444.

- Kirkpatrick, J.F. and A. Turner. 2002. Reversibility of action and safety during pregnancy of immunization against porcine zona pellucida in wild mares (*Equus caballus*). *Reproduction Supplement* 60:197-202.
- Kirkpatrick, J.F. and A. Turner. 2003. Absence of effects from immunocontraception on seasonal birth patterns and foal survival among barrier island wild horses. *Journal of Applied Animal Welfare Science* 6:301-308.
- Kirkpatrick, J.F., A.T. Rutberg, and L. Coates-Markle. 2010. Immunocontraceptive reproductive control utilizing porcine zona pellucida (PZP) in federal wild horse populations, 3rd edition. P.M. Fazio, editor. Downloaded from <http://www.einsten.net/pdf/110242569.pdf>
- Kirkpatrick, J.F., R.O. Lyda, and K. M. Frank. 2011. Contraceptive vaccines for wildlife: a review. *American Journal of Reproductive Immunology* 66:40-50.
- Kirkpatrick, J.F., A.T. Rutberg, L. Coates-Markle, and P.M. Fazio. 2012. Immunocontraceptive Reproductive Control Utilizing Porcine Zona Pellucida (PZP) in Federal Wild Horse Populations. Science and Conservation Center, Billings, Montana.
- Knight, C.M. 2014. The effects of porcine zona pellucida immunocontraception on health and behavior of feral horses (*Equus caballus*). Graduate thesis, Princeton University.
- Levy, J.K., J.A. Friary, L.A. Miller, S.J. Tucker, and K.A. Fagerstone. 2011. Long-term fertility control in female cats with GonaCon™, a GnRH immunocontraceptive. *Theriogenology* 76:1517-1525.
- Liu, I.K.M., M. Bernoco, and M. Feldman. 1989. Contraception in mares heteroimmunized with pig zona pellucida. *Journal of Reproduction and Fertility*, 85:19-29.
- Madosky, J.M., Rubenstein, D.I., Howard, J.J. and Stuska, S., 2010. The effects of immunocontraception on harem fidelity in a feral horse (*Equus caballus*) population. *Applied Animal Behaviour Science*, 128:50-56.
- Magiafoglou, A., M. Schiffer, A.A. Hoffman, and S.W. McKechnie. 2003. Immunocontraception for population control: will resistance evolve? *Immunology and Cell Biology* 81:152-159.
- Mask, T.A., K.A. Schoenecker, A.J. Kane, J.I. Ransom, and J.E. Bruemmer. 2015. Serum antibody immunoreactivity to equine zona protein after SpayVac vaccination. *Theriogenology*, 84:261-267.
- Miller, L.A., J.P. Gionfriddo, K.A. Fagerstone, J.C. Rhyon, and G.J. Killian. 2008. The Single-Shot GnRH Immunocontraceptive Vaccine (GonaCon™) in White-Tailed Deer: Comparison of Several GnRH Preparations. *American Journal of Reproductive Immunology* 60:214-223.

- Miller, L.A., K.A. Fagerstone, and D.C. Eckery. 2013. Twenty years of immunocontraceptive research: lessons learned. *Journal of Zoo and Wildlife Medicine* 44:S84-S96.
- Mills, L.S. and F.W. Allendorf. 1996. The one-migrant-per-generation rule in conservation and management. *Conservation Biology* 10:1509-1518.
- National Park Service (NPS). 2008. Environmental Assessment of Alternatives for Managing the Feral Horses of Assateague Island National Seashore. NPS Assateague Island National Seashore.
- National Research Council of the National Academies of Sciences (NAS). 2013. Using science to improve the BLM wild horse and burro program: a way forward. National Academies Press. Washington, DC.
- Nettles, V. F. 1997. Potential consequences and problems with wildlife contraceptives. *Reproduction, Fertility and Development* 9, 137–143.
- Nolan, M.B., H.J. Bertschinger, and M.L. Schulman. 2018a. Antibody response and safety of a novel recombinant *Zona Pellucida* vaccine formulation in mares. *Journal of Equine Veterinary Science* 66:97.
- Nolan, M.B., H.J. Bertschinger, M. Crampton, and M.L. Schulman. 2018b. Serum anti-Müllerian hormone following *Zona Pellucida* immunocontraceptive vaccination of mares. *Journal of Equine Veterinary Science* 66:105.
- Nolan, M.B., H.J. Bertschinger, R.Roth, M. Crampton, I.S. Martins, G.T. Fosgate, T.A. Stout, and M.L. Schulman. 2018c. Ovarian function following immunocontraceptive vaccination of mares using native porcine and recombinant zona pellucida vaccines formulated with a non-Freund's adjuvant and anti-GnRH vaccines. *Theriogenology* 120:111-116.
- Nuñez, C.M.V., J.S. Adelman, C. Mason, and D.I. Rubenstein. 2009. Immunocontraception decreases group fidelity in a feral horse population during the non-breeding season. *Applied Animal Behaviour Science* 117:74-83.
- Nuñez, C.M., J.S. Adelman, and D.I. Rubenstein. 2010. Immunocontraception in wild horses (*Equus caballus*) extends reproductive cycling beyond the normal breeding season. *PLoS one*, 5(10), p.e13635.
- Nuñez, C.M.V, J.S. Adelman, J. Smith, L.R. Gesquiere, and D.I. Rubenstein. 2014. Linking social environment and stress physiology in feral mares (*Equus caballus*): group transfers elevate fecal cortisol levels. *General and Comparative Endocrinology*. 196:26-33.
- Nuñez, C.M., J.S. Adelman, H.A. Carr, C.M. Alvarez, and D.I. Rubenstein. 2017. Lingering effects of contraception management on feral mare (*Equus caballus*) fertility and social behavior. *Conservation Physiology* 5(1): cox018; doi:10.1093/conphys/cox018.

- Núñez, C.M.V. 2018. Consequences of porcine zona pellucida immunocontraception to feral horses. *Human-Wildlife Interactions* 12:131-142.
- Powell, D.M. 1999. Preliminary evaluation of porcine zona pellucida (PZP) immunocontraception for behavioral effects in feral horses (*Equus caballus*). *Journal of Applied Animal Welfare Science* 2:321-335.
- Powell, D.M. and S.L. Monfort. 2001. Assessment: effects of porcine zona pellucida immunocontraception on estrous cyclicity in feral horses. *Journal of Applied Animal Welfare Science* 4:271-284.
- Powers, J.G., D.L. Baker, T.L. Davis, M.M. Conner, A.H. Lothridge, and T.M. Nett. 2011. Effects of gonadotropin-releasing hormone immunization on reproductive function and behavior in captive female Rocky Mountain elk (*Cervus elaphus nelsoni*). *Biology of Reproduction* 85:1152-1160.
- Powers, J.G., D.L. Baker, M.G. Ackerman, J.E. Bruemmer, T.R. Spraker, M.M. Conner, and T.M. Nett. 2012. Passive transfer of maternal GnRH antibodies does not affect reproductive development in elk (*Cervus elaphus nelson*) calves. *Theriogenology* 78:830-841.
- Powers, J.G., D.L. Baker, R.J. Monello, T.J. Spraker, T.M. Nett, J.P. Gionfriddo, and M.A. Wild. 2013. Effects of gonadotropin-releasing hormone immunization on reproductive function and behavior in captive female Rocky Mountain elk (*Cervus elaphus nelsoni*). *Journal of Zoo and Wildlife Medicine meeting abstracts* S147.
- Ransom, J.I. and B.S. Cade. 2009. Quantifying equid behavior: A research ethogram for free-roaming feral horses. U.S. Geological Survey Techniques and Methods Report 2-A9.
- Ransom, J.I., B.S. Cade, and N.T. Hobbs. 2010. Influences of immunocontraception on time budgets, social behavior, and body condition in feral horses. *Applied Animal Behaviour Science* 124:51-60.
- Ransom, J.I., J.E. Roelle, B.S. Cade, L. Coates-Markle, and A.J. Kane. 2011. Foaling rates in feral horses treated with the immunocontraceptive porcine zona pellucida. *Wildlife Society Bulletin* 35:343-352.
- Ransom, J.I., N.T. Hobbs, and J. Bruemmer. 2013. Contraception can lead to trophic asynchrony between birth pulse and resources. *PLoS one*, 8(1), p.e54972.
- Ransom, J.I., J.G. Powers, N.T. Hobbs, and D.L. Baker. 2014a. Ecological feedbacks can reduce population-level efficacy of wildlife fertility control. *Journal of Applied Ecology* 51:259-269.
- Ransom, J.I., J.G. Powers, H.M. Garbe, M.W. Oehler, T.M. Nett, and D.L. Baker. 2014b. Behavior of feral horses in response to culling and GnRH immunocontraception. *Applied Animal Behaviour Science* 157: 81-92.

- Roelle, J.E., and J.I. Ransom. 2009. Injection-site reactions in wild horses (*Equus caballus*) receiving an immunocontraceptive vaccine: U.S. Geological Survey Scientific Investigations Report 2009–5038.
- Roelle, J.E., F.J. Singer, L.C. Zeigenfuss, J.I. Ransom, F.L. Coates-Markle, and K.A. Schoenecker. 2010. Demography of the Pryor Mountain Wild Horses, 1993-2007. U.S. Geological Survey Scientific Investigations Report 2010–5125.
- Roelle, J.E. and S.J. Oyler-McCance. 2015. Potential demographic and genetic effects of a sterilant applied to wild horse mares. US Geological Survey Open-file Report 2015-1045.
- Roelle, J.E., S.S. Germaine, A.J. Kane, and B.S. Cade. 2017. Efficacy of SpayVac ® as a contraceptive in feral horses. *Wildlife Society Bulletin* 41:107-115.
- Rubenstein, D.I. 1981. Behavioural ecology of island feral horses. *Equine Veterinary Journal* 13:27-34.
- Rutberg, A., K. Grams, J.W. Turner, and H. Hopkins. 2017. Contraceptive efficacy of priming and boosting does of controlled-release PZP in wild horses. *Wildlife Research*: <http://dx.doi.org/10.1071/WR16123>
- Sacco, A.G., M.G. Subramanian, and E.C. Yurewicz. 1981. Passage of zona antibodies via placenta and milk following active immunization of female mice with porcine zonae pellucidae. *Journal of Reproductive Immunology* 3:313-322.
- Sarker, N., M. Tsudzuki, M. Nishibori, and Y. Yamamoto. 1999. Direct and correlated response to divergent selection for serum immunoglobulin M and G levels in chickens. *Poultry Science* 78:1-7.
- Schaut, R.G., M.T. Brewer, J.M. Hostetter, K. Mendoza, J.E. Vela-Ramirez, S.M. Kelly, J.K. Jackman, G. Dell'Anna, J.M. Howard, B. Narasimhan, and W. Zhou. 2018. A single dose polyanhydride-based vaccine platform promotes and maintains anti-GnRH antibody titers. *Vaccine* 36:1016-1023.
- Schulman, M.L., A.E. Botha, S.B. Muenscher, C.H. Annandale, A.J. Guthrie, and H.J. Bertschinger. 2013. Reversibility of the effects of GnRH-vaccination used to suppress reproductive function in mares. *Equine Veterinary Journal* 45:111-113.
- Science and Conservation Center (SCC). 2015. Materials Safety Data Sheet, ZonaStat-H. Billings, Montana.
- Shumake, S.A. and G. Killian. 1997. White-tailed deer activity, contraception, and estrous cycling. Great Plains Wildlife Damage Control Workshop Proceedings, Paper 376.
- Skinner, S.M., Mills, T., Kirchick, H.J. and Dunbar, B.S., 1984. Immunization with Zona Pellucida Proteins Results in Abnormal Ovarian Follicular Differentiation and Inhibition of Gonadotropin-induced Steroid Secretion. *Endocrinology*, 115:2418-2432.

- Stout, T.A.E., J.A. Turkstra, R.H. Meloen, and B. Colenbrander. 2003. The efficacy of GnRH vaccines in controlling reproductive function in horses. Abstract of presentation from symposium, "Managing African elephants: act or let die? Utrecht University, Utrecht, Netherlands.
- Turner, J.W., I.K.M. Liu, and J.F. Kirkpatrick. 1996. Remotely delivered immunocontraception in free-roaming feral burros (*Equus asinus*). *Journal of Reproduction and Fertility* 107:31-35.
- Turner, J.W., I.K. Liu, A.T. Rutberg, and J.F. Kirkpatrick. 1997. Immunocontraception limits foal production in free-roaming feral horses in Nevada. *Journal of Wildlife Management* 61:873-880.
- Turner, J.W., I.K. Liu, D.R. Flanagan, K.S. Bynum, and A.T. Rutberg. 2002. Porcine zona pellucida (PZP) immunocontraception of wild horses (*Equus caballus*) in Nevada: a 10 year study. *Reproduction Supplement* 60:177-186.
- Turner, J.W., and J.F. Kirkpatrick. 2002. Effects of immunocontraception on population, longevity and body condition in wild mares (*Equus caballus*). *Reproduction (Cambridge, England) Supplement*, 60, pp.187-195.
- Turner, J.W., I.K. Liu, D.R. Flanagan, A.T. Rutberg, and J.F. Kirkpatrick. 2007. Immunocontraception in wild horses: one inoculation provides two years of infertility. *Journal of Wildlife Management* 71:662-667.
- Turner, J.W., A.T. Rutberg, R.E. Naugle, M.A. Kaur, D.R. Flanagan, H.J. Bertschinger, and I.K.M. Liu. 2008. Controlled-release components of PZP contraceptive vaccine extend duration of infertility. *Wildlife Research* 35:555-562.
- US Fish and Wildlife Service (USFWS). 2015. Endangered and Threatened Wildlife and Plants; 90-day findings on 31 petitions. *Federal Register* 80 (126):37568-37579.
- Wang-Cahill, F., J. Warren, T. Hall, J. O'Hare, A. Lemay, E. Ruell, and R. Wimberly. In press. Use of GonaCon in wildlife management. Chapter 24 in USDA-APHIS, Human health and ecological risk assessment for the use of wildlife damage management methods by APHIS-Wildlife Services. USDA APHIS, Fort Collins, Colorado.
- Wright, S. 1931. Evolution in Mendelian populations. *Genetics* 16:97-159.
- Yao, Z., W. Si, W. Tian, J. Ye, R. Zhu, X. Li, S. Ki, Q. Zheng, Y. Liu, and F. Fang. 2018. Effect of active immunization using a novel GnRH vaccine on reproductive function in rats. *Theriogenology* 111:1-8. <https://doi.org/10.1016/j.theriogenology.2018.01.013>
- Zoo Montana. 2000. Wildlife Fertility Control: Fact and Fancy. Zoo Montana Science and Conservation Biology Program, Billings, Montana.

G.4. Effects of Sterilization, Including Spaying and Neutering

Various forms of fertility control can be used in wild horses and wild burros, with the goals of maintaining herds at or near AML, reducing fertility rates, and reducing the frequency of gathers and removals. The WFRHBA of 1971 specifically provides for contraception and sterilization (16 U.S.C. 1333 section 3.b.1). Fertility control measures have been shown to be a cost-effective and humane treatment to slow increases in wild horse populations or, when used in combination with gathers, to reduce horse population size (Bartholow 2004, de Seve and Boyles-Griffin 2013, Fonner and Bohara 2017). Population growth suppression becomes less expensive if fertility control is long-lasting (Hobbs et al. 2000), such as with sterilization methods that may include spaying and neutering. Sterilizing a female horse (mare) or burro (jenny) can be accomplished by several methods, some of which are surgical and others of which are non-surgical. In this review, ‘spaying’ is defined to be surgical sterilization, usually accomplished by removal of the ovaries, but other surgical methods such as tubal ligation that lead to sterility may also be considered a form of spaying. Unlike in dog and cat spaying, spaying a horse or burro does not entail removal of the uterus. Here, ‘neutering’ is defined to be the sterilization of a male horse (stallion) or burro (jack), either by removal of the testicles (castration, also known as gelding) or by vasectomy, where the testicles are retained but no sperm leave the body by severing or blocking the vas deferens or epididymis.

In the context of BLM wild horse and burro management, sterilization is expected to be successful to the extent that it reduces the number of reproducing females. By definition, sterilizing a given female is 100% effective as a fertility control method for that female. Neutering males may be effective in one of two ways. First, neutered males may continue to guard fertile females, preventing the females from breeding with fertile males. Second, if neutered males are included in a herd that has a high male-to-female sex ratio, then the neutered males may comprise some of the animals within the appropriate management level (AML) of that herd, which would effectively reduce the number of females in the herd. Although these and other fertility control treatments may be associated with a number of potential physiological, behavioral, demographic, and genetic effects, those impacts are generally minor and transient, do not prevent overall maintenance of a self-sustaining population, and do not generally outweigh the potential benefits of using contraceptive treatments in situations where it is a management goal to reduce population growth rates (Garrott and Oli 2013).

Peer-reviewed scientific literature details the expected impacts of sterilization methods on wild horses and burros. No finding of excess animals is required for BLM to pursue sterilization in wild horses or wild burros, but NEPA analysis has been required. This review focuses on peer-reviewed scientific literature. The summary that follows first examines effects of female sterilization, then neuter use in males. This review does not examine effects of reversible fertility control vaccines. Cited studies are generally limited to those involving horses and burros, except where including studies on other species helps in making inferences about physiological or behavioral questions not yet addressed in horses or burros specifically. While most studies reviewed here refer to horses, burros are extremely similar in terms of physiology, such that expected effects are comparable, except where differences between the species are noted.

On the whole, the identified impacts at the herd level are generally transient. The principle impact to individuals treated is sterility, which is the intended outcome. Sterilization that affects individual horses and burros does not prevent BLM from ensuring that there will be self-sustaining populations of wild horses and burros in single HMAs, in complexes of HMAs, and at regional scales of multiple HMAs and complexes. Under the WFRHBA of 1971, BLM is charged with maintaining self-reproducing populations of wild horses and burros. The National Academies of Sciences (2013) encouraged BLM to manage wild horses and burros at the spatial scale of “metapopulations” – that is, across multiple HMAs and complexes in a region. In fact, many HMAs have historical and ongoing genetic and demographic connections with other HMAs, and BLM routinely moves animals from one to another to improve local herd traits and maintain high genetic diversity.

Discussions about herds that are ‘non-reproducing’ in whole or in part are in the context of this ‘metapopulation’ structure, where self-sustaining herds are not necessarily at the scale of single HMAs. So long as the definition of what constitutes a self-sustaining herd includes the larger set of HMAs that have past or ongoing demographic and genetic connections – as is recommended by the NAS 2013 report – it is clear that single HMAs can be managed as non-reproducing in whole or in part while still allowing for a self-sustaining population of wild horses or burros at the broader spatial scale. Wild horses are not an endangered species (USFWS 2015), nor are they rare. Nearly 67,000 adult wild horses and nearly 15,000 adult wild burros roam BLM lands as of March 1, 2018, and those numbers do not include at least 10,000 WH&B on US Forest Service lands, and at least 50,000 feral horses on tribal lands in the Western United States.

All fertility control methods affect the behavior and physiology of treated animals (NAS 2013), and are associated with potential risks and benefits, including effects of handling, frequency of handling, physiological effects, behavioral effects, and reduced population growth rates (Hampton et al. 2015). Contraception methods alone do not remove excess horses from an HMA’s population, so one or more gathers are usually needed to bring the herd down to a level close to AML. Horses are long-lived, potentially reaching 20 years of age or more in the wild. Except in cases where extremely high fractions of mares are rendered infertile over long time periods of (i.e., 10 or more years), spaying and neutering are not very effective at reducing population growth rates to the point where births equal deaths in a herd. However, even modest levels of fertility control activities can reduce the frequency of horse gather activities, and costs to taxpayers. Population growth suppression becomes less expensive if fertility control is long-lasting (Hobbs et al. 2000), such as with sterilization. Because sterilizing animals requires capturing and handling, the risks and costs associated with capture and handling of horses may be comparable to those of gathering for removal, but with expectedly lower adoption and long-term holding costs.

Effects of handling and marking

Surgical sterilization techniques, while not reversible, may control horse reproduction without the kind of additional handling or darting that can be needed to administer contraceptive vaccines. In this sense, sterilization surgeries can be used to achieve herd management objectives with a relative minimum level of animal handling and management over the long term. The WFRHBA (as amended) indicates that management should be at the minimum level necessary to achieve management objectives (CFR 4710.4), and if gelding some fraction of a managed

population can reduce population growth rates by replacing breeding mares, it then follows that spaying or neutering some individuals can lead to a reduced number of handling occasions and removals of excess horses from the range, which is consistent with legal guidelines. Other fertility control options that may be temporarily effective on male horses, such as the injection of GonaCon-Equine immunocontraceptive vaccine, apparently require multiple handling occasions to achieve longer-term male infertility. Similarly, some formulations of PZP immunocontraception that is currently available for use in female wild horses and burros require handling or darting every year (though longer-term effects may result after 4 or more treatments; Nuñez et al. 2017). By some measures, any management activities that require multiple capture operations to treat a given individual would be more intrusive for wild horses and potentially less sustainable than an activity that requires only one handling occasion.

It is prudent for sterilized animals to be readily identifiable, either via freeze brand marks or unique coloration, so that their treatment history is easily recognized (e.g., BLM 2010). Markings may also be useful into the future to determine the approximate fraction of geldings in a herd and could provide additional insight regarding gather efficiency. BLM has instituted capture and animal welfare program guidelines to reduce the sources of handling stress in captured animals (BLM 2015). Handling may include freeze-marking, for the purpose of identifying an individual. Some level of transient stress is likely to result in newly captured horses that are not previously marked. Under past management practices, captured horses experienced increased, transient stress levels from handling (Ashley and Holcombe 2001). It is difficult to compare that level of temporary stress with long-term stress that can result from food and water limitation on the range (e.g., Creel et al. 2013), which could occur in the absence of herd management.

Most horses recover from the stress of capture and handling quickly once released back to the range, and none are expected to suffer serious long-term effects from gelding, other than the direct consequence of becoming infertile.

Observations of the long-term outcomes of sterilization may be recorded during routine resource monitoring work. Such observations could include but not be limited to band size, social interactions with other geldings and harem bands, distribution within their habitat, forage utilization and activities around key water sources. Periodic population inventories and future gather statistics could provide additional anecdotal information.

G.4.1. Neutering Males

Castration (the surgical removal of the testicles, also called gelding or neutering) is a surgical procedure for the horse sterilization that has been used for millennia. Vasectomy involves severing or blocking the vas deferens or epididymis, to prevent sperm from being ejaculated. The procedures are fairly straight forward and has a relatively low complication rate. As noted in the review of scientific literature that follows, the expected effects of gelding and vasectomy are well understood overall, even though there is some degree of uncertainty about the exact quantitative outcomes for any given individual (as is true for any natural system).

Including a portion of neutered males in a herd can lead to a reduced population-level per-capita growth rate if they cause a marginal decrease in female fertility or if the neutered males take

some of the places that would otherwise be occupied by fertile females. By having a skewed sex ratio with fewer females than males (fertile stallions plus neutered males), the result will be that there will be a lower number of breeding females in the population. Including neutered males in herd management is not new for BLM and federal land management. Geldings have been released on BLM lands as a part of herd management in the Barren Valley complex in Oregon (BLM 2011), the Challis HMA in Idaho (BLM 2012), and the Conger HMA in Utah (BLM 2016). Vasectomized males and geldings were also included in US Fish and Wildlife Service management plans for the Sheldon National Wildlife Refuge that relied on sterilization and removals (Collins and Kasbohm 2016). Taking into consideration the literature available at the time, the National Academies of Sciences concluded in their 2013 report that a form of vasectomy was one of the three most promising methods for WH&B fertility control (NAS 2013).

Nelson (1980) and Garrott and Siniff (1992) modeled potential efficacy of male-oriented contraception as a population management tool, and both studies agreed that while slowing growth, sterilizing only dominant males (i.e., harem-holding stallions) would result in only marginal reduction in female fertility rates. Eagle et al. (1993) and Asa (1999) tested this hypothesis on HMAs where dominant males were vasectomized. Their findings agreed with modeling results from previous studies, and they also concluded that sterilizing only dominant males would not provide the desired reduction in female fertility and overall population growth rate, assuming that the numbers of fertile females is not changed. While bands with vasectomized harem stallions tended to have fewer foals, breeding by bachelors and subordinate stallions meant that population growth still occurred – female fertility was not dramatically reduced. Collins and Kasbohm (2016) demonstrated that there was a reduced fertility rate in a feral horse herd with both spayed and vasectomized horses – some geldings were also present in that herd. Garrott and Siniff (1992) concluded from their modeling that male sterilization would effectively cause there to be zero population growth (the point where births roughly equal deaths) only if a large proportion of males (i.e., >85%) could be sterilized. In cases where the goal of harem stallion sterilization is to reduce population growth rates, success appears to be dependent on a stable group structure, as strong bonds between a stallion and mares reduce the probability of a mare mating an extra-group stallion (Nelson 1980, Garrott and Siniff 1992, Eagle et al. 1993, Asa 1999).

Despite these studies, neutered males can be used to reduce overall growth rates in a management strategy that does not rely on any expectation that geldings will retain harems or lead to a reduction in per-female fertility rates. The primary goal of including neutered males in a herd need not necessarily be to reduce female fertility (although that may be one result). Rather, by including some neutered males in a herd that also has fertile mares and stallions, the neutered males would take some of the spaces toward AML that would otherwise be taken by fertile females. If the total number of horses is constant but neutered males are included in the herd, this can reduce the number of fertile mares, therefore reducing the absolute number of foals produced. Put another way, if neutered males occupy spaces toward AML that would otherwise be filled by fertile mares, that will reduce growth rates merely by the fact of causing there to be a lower starting number of fertile mares.

Direct Effects of Neutering

No animals which appear to be distressed, injured, or in poor health or condition would be selected for gelding. Stallions would not typically be neutered within 72 hours of capture. The surgery would be performed by a veterinarian using general anesthesia and appropriate surgical techniques. The final determination of which specific animals would be gelded would be based on the professional opinion of the attending veterinarian in consultation with the Authorized Officer (i.e., See the SOPs for neutering in the Antelope / Triple B gather EA, DOI-BLM-NV-E030-2017-010-EA).

Though neutering males is a common surgical procedure, especially gelding, some level of minor complications after surgery may be expected (Getman 2009), and it is not always possible to predict when postoperative complications would occur. Fortunately, the most common complications are almost always self-limiting, resolving with time and exercise. Individual impacts to the stallions during and following the gelding process should be minimal and would mostly involve localized swelling and bleeding. Complications may include, but are not limited to: minor bleeding, swelling, inflammation, edema, infection, peritonitis, hydrocele, penile damage, excessive hemorrhage, and eventration (Schumacher 1996, Searle et al. 1999, Getman 2009). A small amount of bleeding is normal and generally subsides quickly, within 2-4 hours following the procedure. Some degree of swelling is normal, including swelling of the prepuce and scrotum, usually peaking between 3-6 days after surgery (Searle et al. 1999). Swelling should be minimized through the daily movements (exercise) of the horse during travel to and from foraging and watering areas. Most cases of minor swelling should be back to normal within 5-7 days, more serious cases of moderate to severe swelling are also self-limiting and are expected to resolve with exercise after one to 2 weeks. Older horses are reported to be at greater risk of post-operative edema, but daily exercise can prevent premature closure of the incision and prevent fluid buildup (Getman 2009). In some cases, a hydrocele (accumulation of sterile fluid) may develop over months or years (Searle et al. 1999). Serious complications (eventration, anesthetic reaction, injuries during handling, etc.) that result in euthanasia or mortality during and following surgery are rare (e.g., eventration rate of 0.2% to 2.6% noted in Getman 2009, but eventration rate of 4.8% noted in Shoemaker et al. 2004) and vary according to the population of horses being treated (Getman 2009). Normally one would expect serious complications in less than 5% of horses operated under general anesthesia, but in some populations these rates have been as high as 12% (Shoemaker 2004). Serious complications are generally noted within 3 or 4 hours of surgery but may occur any time within the first week following surgery (Searle et al. 1999). If they occur, they would be treated with surgical intervention when possible, or with euthanasia when there is a poor prognosis for recovery. Vasectomized stallions may remain fertile for up to 6 weeks after surgery, so it is optimal if that treatment occurs well in advance of the season of mare fertility starting in the spring (NAS 2013). The NAS report (2013) suggested that chemical vasectomy, which has been developed for dogs and cats, may be appropriate for wild horses and burros.

For intact stallions, testosterone levels appear to vary as a function of age, season, and harem size (Khalil et al 1998). It is expected that testosterone levels will decline over time after castration. Testosterone levels should not change due to vasectomy. Vasectomized stallions should retain their previous levels of libido. Domestic geldings had a significant prolactin response to sexual stimulation but lacked the cortisol response present in stallions (Colborn et al.

1991). Although libido and the ability to ejaculate tends to be gradually lost after castration (Thompson et al. 1980), some geldings continue to mount mares and intromit (Rios and Houpt 1995, Schumacher 2006).

Indirect Effects of Neutering

Other than the short-term outcomes of surgery, neutering is not expected to reduce males' survival rates. Castration is actually thought to increase survival as males are released from the cost of reproduction (Jewell 1997). In Soay sheep castrates survived longer than rams in the same cohort (Jewell 1997), and Misaki horse geldings lived longer than intact males (Kaseda et al. 1997, Khalil and Murakami 1999). Moreover, it is unlikely that a reduced testosterone level will compromise gelding survival in the wild, considering that wild mares survive with low levels of testosterone. Consistent with geldings not expending as much energy toward in attempts to obtain or defend a harem, it is expected that wild geldings may have a better body condition than wild, fertile stallions. In contrast, vasectomized males may continue to defend or compete for harems in the way that fertile males do, so they are not expected to experience an increase in health or body condition due to surgery.

Depending on whether an HMA is non-reproducing in whole or in part, reproductive stallions may or may not still be a component of the population's age and sex structure. The question of whether a given neutered male would or would not attempt to maintain a harem is not germane to population-level management. It is worth noting, though, that the BLM is not required to manage populations of wild horses in a manner that ensures that any given individual maintains its social standing within any given harem or band. Neutering a subset of stallions would not prevent other fertile stallions and mares from continuing with the typical range of social behaviors for sexually active adults. For fertility control strategies where gelding is intended to reduce growth rates by virtue of sterile males defending harems, the NAS (2013) suggested that the effectiveness of gelding on overall reproductive rates may depend on the pre-castration social roles of those animals. Having a post-gather herd with some neutered males and a lower fraction of fertile mares necessarily reduces the absolute number of foals born per year, compared to a herd that includes more fertile mares. An additional benefit is that geldings that would otherwise be permanently removed from the range (for adoption, sale or other disposition) may be released back onto the range where they can engage in free-roaming behaviors.

Behavioral Effects of Neutering

Feral horses typically form bands composed of an adult male with 1 to 3 adult females and their immature offspring (Feist and McCullough 1976, Berger 1986, Roelle et al. 2010). In many populations subordinate 'satellite' stallions have been observed associating with the band, although the function of these males continues to be debated (see Feh 1999, and Linklater and Cameron 2000). Juvenile offspring of both sexes leave the band at sexual maturity (normally around two or three years of age (Berger 1986), but adult females may remain with the same band over a span of years. Group stability and cohesion is maintained through positive social interactions and agonistic behaviors among all members, and herding and reproductive behaviors from the stallion (Ransom and Cade 2009). Group movements and consortship of a stallion with mares is advertised to other males through the group stallion marking dung piles as they are encountered, and over-marking mare eliminations as they occur (King and Gurnell 2006).

In horses, males play a variety of roles during their lives (Deniston 1979): after dispersal from their natal band they generally live as bachelors with other young males, before associating with mares and developing their own breeding group as a harem stallion or satellite stallion. In any population of horses not all males will achieve harem stallion status, so all males do not have an equal chance of breeding (Asa 1999). Stallion behavior is thought to be related to androgen levels, with breeding stallions having higher androgen concentrations than bachelors (Angle et al. 1979, Chaudhuri and Ginsberg 1990, Khalil et al. 1998). A bachelor with low libido had lower levels of androgens, and two-year-old bachelors had higher testosterone levels than two year olds with undescended testicles who remained with their natal band (Angle et al. 1979).

Vasectomized males continue to attempt to defend or gain breeding access to females. It is generally expected that vasectomized WH&B will continue to behave like fertile males, given that the only physiological change in their condition is a lack of sperm in their ejaculate. If a vasectomized stallion retains a harem, the females in the harem will continue to cycle until they are fertilized by another stallion, or until the end of the breeding season. As a result, the vasectomized stallion may be involved in more aggressive behaviors to other males through the entire breeding season (Asa 1999), which may divert time from foraging and cause him to be in poorer body condition going into winter. Ultimately, this may lead to the stallion losing control of a given harem. A feral horse herd with high numbers of vasectomized stallions retained typical harem social structure (Collins and Kasbohm 2016). Again, it is worth noting that the BLM is not required to manage populations of wild horses in a manner that ensures that any given individual maintains its social standing within any given harem or band.

Neutering males by gelding adult male horses is expected to result in reduced testosterone production, which is expected to directly influence reproductive behaviors (NAS 2013). However, testosterone levels alone are not a predictor of masculine behavior (Line et al. 1985, Schumacher 2006). In domestic geldings, 20-30% continued to show stallion-like behavior, whether castrated pre- or post-puberty (Line et al. 1985). Gelding of domestic horses most commonly takes place before or shortly after sexual maturity, and age-at-gelding can affect the degree to which stallion-like behavior is expressed later in life. In intact stallions, testosterone levels peak increase up to an age of ~4-6 years, and can be higher in harem stallions than bachelors (Khalil et al 1998). It is assumed that free roaming wild horse geldings would generally exhibit reduced aggression toward other horses and reduced reproductive behaviors (NAS 2013). The behavior of wild horse geldings in the presence of intact stallions has not been well documented, but the literature review below can be used to make reasonable inferences about their likely behaviors.

Despite livestock being managed by neutering males for millennia, there is relatively little published research on castrates' behaviors (Hart and Jones 1975). Stallion behaviors in wild or pasture settings are better documented than gelding behaviors, but inferences about how the behaviors of geldings will change, how quickly any change will occur after surgery, or what effect gelding an adult stallion and releasing him back in to a wild horse population will have on his behavior and that of the wider population must be surmised from the existing literature. There is an ongoing BLM study in Utah focused on the individual and population-level effects of including some geldings in a free-roaming horse population (BLM 2016) but results from that

study are not yet available. However, inferences about likely behavioral outcomes of gelding can be made based on available literature.

The effect of castration on aggression in horses has not often been quantified. One report has noted that high levels of aggression continued to be observed in domestic horse geldings who also exhibited sexual behaviors (Rios and Houpt 1995). Stallion-like behavior in domestic horse geldings is relatively common (Smith 1974, Schumacher 1996), being shown in 20-33% of cases whether the horse was castrated pre- or post-puberty (Line et al. 1985, Rios and Houpt 1995, Schumacher 2006). While some of these cases may be due to cryptorchidism or incomplete surgery, it appears that horses are less dependent on hormones than other mechanisms for the maintenance of sexual behaviors (Smith 1974). Domestic geldings exhibiting masculine behavior had no difference in testosterone concentrations than other geldings (Line et al. 1985, Schumacher 2006), and in some instances the behavior appeared context dependent (Borsberry 1980, Pearce 1980).

Dogs and cats are commonly neutered, and it is also common for them to continue to exhibit reproductive behaviors several years after castration (Dunbar 1975). Dogs, ferrets, hamsters, and marmosets continued to show sexually motivated behaviors after castration, regardless of whether they had previous experience or not, although in beagles and ferrets there was a reduction in motivation post-operatively (Hart 1968, Dunbar 1975, Dixson 1993, Costantini et al. 2007, Vinke et al. 2008). Ungulates continued to show reproductive behaviors after castration, with goats and llamas continuing to respond to females even a year later in the case of goats, although mating time and the ejaculatory response was reduced (Hart and Jones 1975, Nickolmann et al. 2008).

The likely effects of castration on geldings' social interactions and group membership can be inferred from available literature. In a pasture study of domestic horses, Van Dierendonk et al. (1995) found that social rank among geldings was directly correlated to the age at which the horse was castrated, suggesting that social experiences prior to sterilization may influence behavior afterward. Of the two geldings present in a study of semi-feral horses in England, one was dominant over the mares whereas a younger gelding was subordinate to older mares; stallions were only present in this population during a short breeding season (Tyler 1972). A study of domestic geldings in Iceland held in a large pasture with mares and sub-adults of both sexes, but no mature stallions, found that geldings and sub-adults formed associations amongst each other that included interactions such as allo-grooming and play, and were defined by close proximity (Sigurjónsdóttir et al. 2003). These geldings and sub-adults tended to remain in a separate group from mares with foals, similar to castrated Soay sheep rams (*Ovis aries*) behaving like bachelors and grouping together or remaining in their mother's group (Jewell 1997). In Japan, Kaseda et al. (1997) reported that young males dispersing from their natal harem and geldings moved to a different area than stallions and mares during the non-breeding season. Although the situation in Japan may be the equivalent of a bachelor group in natural populations, in Iceland this division between mares and the rest of the horses in the herd contradicts the dynamics typically observed in a population containing mature stallions. Sigurjónsdóttir et al. (2003) also noted that in the absence of a stallion, allo-grooming between adult females increased drastically. Other findings included increased social interaction among yearlings, display of stallion-like behaviors such as mounting by the adult females, and decreased

association between females and their yearling offspring (Sigurjónsdóttir et al. 2003). In the same population in Iceland Van Dierendonck et al. (2004) concluded that the presence of geldings did not appear to affect the social behavior of mares or negatively influence parturition, mare-foal bonding, or subsequent maternal activities. Additionally, the welfare of broodmares and their foals was not affected by the presence of geldings in the herd (Van Dierendonck et al. 2004). These findings are important because treated geldings will be returned to the range in the presence of pregnant mares and mares with foals of the year.

The likely effects of castration on geldings' home range and habitat use can also be surmised from available literature. Bands of horses tend to have distinct home ranges, varying in size depending on the habitat and varying by season, but always including a water source, forage, and places where horses can shelter from inclement weather or insects (King and Gurnell 2005). By comparison, bachelor groups tend to be more transient, and can potentially use areas of good forage further from water sources, as they are not constrained by the needs of lactating mares in a group. The number of observations of gelded wild stallion behavior are still too few to make general predictions about whether a particular gelded stallion individual will behave like a harem stallion, a bachelor, or form a group with geldings that may forage and water differently from fertile wild horses.

Sterilizing wild horses does not change their status as wild horses under the WFRHBA (as amended). In terms of whether geldings will continue to exhibit the free-roaming behavior that defines wild horses, BLM does expect that geldings would continue to roam unhindered once they are returned to the range. Wild horse movements may be motivated by a number of biological impulses, including the search for forage, water, and social companionship that is not of a sexual nature. As such, a gelded animal would still be expected to have a number of internal reasons for moving across a landscape and, therefore, exhibiting 'free-roaming' behavior. Despite marginal uncertainty about subtle aspects of potential changes in habitat preference, there is no expectation that gelding wild horses will cause them to lose their free-roaming nature. It is worth noting that individual choices in wild horse group membership, home range, and habitat use are not protected under the WFRHBA. BLM acknowledges that geldings may exhibit some behavioral differences after surgery, compared to intact stallions, but those differences are not be expected to remove the geldings' rebellious and feisty nature, or their defiance of man. While it may be that a gelded horse could have a different set of behavioral priorities than an intact stallion, the expectation is that geldings will choose to act upon their behavioral priorities in an unhindered way, just as is the case for an intact stallion. In this sense, a gelded male would be just as much 'wild' as defined by the WFRHBA as any intact stallion, even if his patterns of movement differ from those of an intact stallion. Congress specified that sterilization is an acceptable management action (16 USC § 1333.b.1). Sterilization is not one of the clearly defined events that cause an animal to lose its status as a wild free-roaming horse (16 USC § 1333.2.C.d). Several academics have offered their opinions about whether gelding a given stallion would lead to that individual effectively losing its status as a wild horse (Rutberg 2011, Kirkpatrick 2012, Nock 2017). Those opinions are based on a semantic and subjective definition of 'wild,' while BLM must adhere to the legal definition of what constitutes a wild horse, based on the WFRHBA (as amended). Those individuals have not conducted any studies that would test the speculative opinion that gelding wild stallions will cause them to become docile. BLM is not obliged to base management decisions on such opinions, which do not meet the BLM's

principle and practice to “Use the best available scientific knowledge relevant to the problem or decision being addressed, relying on peer reviewed literature when it exists” (Kitchell et al. 2015).

G.4.2. Mare Sterilization

Surgical sterilization (spaying mares by removing a mare’s ovaries), via colpotomy, has been an established veterinary technique since 1903 (Loesch and Rodgerson 2003, NAS 2013). Spaying via colpotomy has the advantage of not leaving any external wound that could become infected. For this reason, it has been identified as a good choice for sterilization of feral or wild mares (Rowland et al. 2018). The procedure has a relatively low complication rate, although post-surgical mortality and morbidity are possible, as with any surgery. Herd-level birth rate is expected to decline in direct proportion to the fraction of spayed mares in the herd because spayed mares cannot become pregnant. Spaying mares has already been shown to be an effective part of feral horse management that reduced herd growth rates on federal lands (Collins and Kasbohm 2016).

Current Methods of Sterilization

This literature review of mare sterilization impacts focuses on 4 methods: spaying via flank laparoscopy, spaying via colpotomy, non-surgical physical sterilization, and pharmacological or immunocontraceptive sterilization. The anticipated effects are both physical and behavioral. Physical effects of surgical methods would be due to post-treatment healing and the possibility for complications.

Colpotomy is a surgical technique in which there is no external incision, reducing susceptibility to infection. For this reason, ovariectomy via colpotomy has been identified as a good choice for feral or wild horses (Rowland et al. 2018). Ovariectomy via colpotomy is a relatively short surgery, with a relatively quick expected recovery time. In 1903, Williams first described a vaginal approach, or colpotomy, using an ecraseur to ovariectomize mares (Loesch and Rodgerson 2003). The ovariectomy via colpotomy procedure has been conducted for over 100 years, normally on open (non-pregnant), domestic mares. It is expected that the surgeon should be able to access ovaries with ease in mares that are in the early- or mid-stage of pregnancy. The anticipated risks associated with the pregnancy are described below. When wild horses are gathered or trapped for fertility control treatment there would likely be mares in various stages of gestation. Removal of the ovaries is permanent and 100 percent effective, however the procedure is not without risk.

Flank laparoscopy (Lee and Hendrickson 2008, Devick et al. 2018, Easley et al. 2018) is commonly used in domestic horses for application in mares due to its minimal invasiveness and full observation of the operative field. Ovariectomy via flank laparoscopy was seen as the lowest risk method considered by a panel of expert reviewers convened by USGS (Bowen 2015). In a review of unilateral and bilateral laparoscopic ovariectomy on 157 mares, Röcken et al. (2011) found that 10.8% of mares had minor post-surgical complications and recorded no mortality. Mortality due to this type of surgery, or post-surgical complications, is not expected, but is a possibility. In two studies, ovariectomy by laparoscopy or endoscope-assisted colpotomy did not cause mares to lose weight, and there was no need for rescue analgesia following surgery (Pader et al. 2011, Bertin et al. 2013). This surgical approach entails three small incisions on the

animal's flank, through which three cannulae (tubes) allow entry of narrow devices to enter the body cavity: these are the insufflator, endoscope, and surgical instrument. The surgical procedure involves the use of narrow instruments introduced into the abdomen via cannulas for the purpose of transecting or sealing (Easley 2018) the ovarian pedicle, but the insufflation should allow the veterinarian to navigate inside the abdomen without damaging other internal organs. The insufflator blows air into the cavity to increase the operating space between organs, and the endoscope provides a video feed to visualize the operation of the surgical instrument. This procedure can require a relatively long duration of surgery but tends to lead to the lowest post-operative rates of complications. Flank laparoscopy may leave three small (<5 cm) visible scars on one side of the horse's flank, but even in performance horses these scars are considered minimal. It is expected that the tissues and musculature under the skin at the site of the incisions in the flank will heal quickly, leaving no long-lasting effects on horse health. Monitoring for up to two weeks at the facility where surgeries take place will allow for veterinary inspection of wound healing. The ovaries may be dropped into the abdomen, but this is not expected to cause any health problem; it is usually done in ovariectomies in cattle (e.g., the Willis Dropped Ovary Technique) and Shoemaker et al. (2014) found no problems with revascularization or necrosis in a study of young horses using this method.

Non-surgical, physical sterilization would include any physical form of sterilization that does not involve surgery. This could include any form of physical procedure that leads a mare to be unable to become pregnant, or to maintain a pregnancy. For example, one form of physical, non-surgical sterilization causes a long-term blockage of the oviduct, so that fertile eggs cannot go from the ovaries to the uterus. One form of this procedure infuses medical cyanoacrylate glue into the oviduct to cause long-term blockage (Bigolin et al. 2009). Treated mares would need to be screened to ensure they are not pregnant. The procedure is transcervical, so the treated mare cannot have a fetus in the uterus at the time of treatment. The mare would be sterile, although she would continue to have estrus cycles.

Pharmacological or immunocontraceptive sterilization methods would use an as-yet undetermined drug or vaccine to cause sterilization. At this time, BLM has not yet identified a pharmacological or immunocontraceptive method to sterilize mares that has been proven to reliably and humanely sterilize wild horse mares. However, there is the possibility that future development and testing of new methods could make an injectable sterilant available for wild horse mares. Analyses of the effects of having sterile mares as a part of a wild horse herd, such as due to surgical sterilization, would likely be applicable to non-surgical methods as well.

Effects of Spaying on Pregnancy and Foal

The average mare gestation period ranges from 335 to 340 days (Evans et al. 1977, p. 373). There are few peer reviewed studies documenting the effects of ovariectomy on the success of pregnancy in a mare. A National Research Council of the National Academies of Sciences (NAS) committee that reviewed research proposals in 2015 explained, "The mare's ovaries and their production of progesterone are required during the first 70 days of pregnancy to maintain the pregnancy" (NAS 2015). In female mammals, less progesterone is produced when ovaries are removed, but production does not cease (Webley and Johnson 1982). In 1977, Evans et al. stated that by 200 days, the secretion of progesterone by the corpora lutea is insignificant because removal of the ovaries does not result in abortion (p. 376). "If this procedure were

performed in the first 120 days of pregnancy, the fetus would be resorbed or aborted by the mother. If performed after 120 days, the pregnancy should be maintained. The effect of ovary removal on a pregnancy at 90–120 days of gestation is unpredictable because it is during this stage of gestation that the transition from corpus luteum to placental support typically occurs” (NAS 2015). In 1979, Holtan et al. evaluated the effects of bilateral ovariectomy at selected times between 25 and 210 days of gestation on 50 mature pony mares. Their results show that abortion (resorption) of the conceptus (fetus) occurred in all 14 mares ovariectomized before day 50 of gestation, that pregnancy was maintained in 11 of 20 mares after ovariectomy between days 50 and 70, and that pregnancy was not interrupted in any of 12 mares ovariectomized on days 140 to 210. Those results are similar to the suggestions of the NAS committee (2015).

For those pregnancies that are maintained following the procedure, likely those past approximately 120 days, the development of the foal is not expected to be affected. However, because this procedure is not commonly conducted on pregnant mares the rate of complications to the fetus has not yet been quantified. There is the possibility that entry to the abdominal cavity could cause premature births related to inflammation. However, after five months the placenta should hormonally support the pregnancy regardless of the presence or absence of ovaries. Gestation length was similar between ovariectomized and control mares (Holtan et al. 1979).

Direct Effects of Spaying

Between 2009 and 2011, the Sheldon NWR in Nevada conducted ovariectomy via colpotomy surgeries (August through October) on 114 feral mares and released them back to the range with a mixture of sterilized stallions and untreated mares and stallions (Collins and Kasbohm 2016). Gestational stage was not recorded, but a majority of the mares were pregnant (Gail Collins, US Fish and Wildlife Service (USFWS), pers. comm.). Only a small number of mares were very close to full term. Those mares with late term pregnancies did not receive surgery as the veterinarian could not get good access to the ovaries due to the position of the foal (Gail Collins, USFWS, pers. comm.). After holding the mares for an average of 8 days after surgery for observation, they were returned to the range with other treated and untreated mares and stallions (Collins and Kasbohm 2016). During holding the only complications were observed within 2 days of surgery. The observed mortality rate for ovariectomized mares following the procedure was less than 2 percent (Collins and Kasbohm 2016, Pielstick pers. comm.).

During the Sheldon NWR ovariectomy study, mares generally walked out of the chute and started to eat; some would raise their tail and act as if they were defecating; however, in most mares one could not notice signs of discomfort (Bowen 2015). In their discussion of ovariectomy via colpotomy, McKinnon and Vasey (2007) considered the procedure safe and efficacious in many instances, able to be performed expediently by personnel experienced with examination of the female reproductive tract, and associated with a complication rate that is similar to or less than male castration. Nevertheless, all surgery is associated with some risk. Loesch et al. (2003) lists that following potential risks with colpotomy: pain and discomfort; injuries to the cervix, bladder, or a segment of bowel; delayed vaginal healing; eventration of the bowel; incisional site hematoma; intraabdominal adhesions to the vagina; and chronic lumbar or bilateral hind limb pain. Most horses, however, tolerate ovariectomy via colpotomy with very few complications, including feral horses (Collins and Kasbohm 2016). Evisceration is also a possibility, but these complications are considered rare (Prado and Schumacher, 2017). Mortality due to surgery or

post-surgical complications is not anticipated, but it is a possibility and therefore every effort would be made to mitigate risks.

In September 2015, the BLM solicited the USGS to convene a panel of veterinary experts to assess the relative merits and drawbacks of several surgical ovariectomy techniques that are commonly used in domestic horses for potential application in wild horses. A table summarizing the various methods was sent to the BLM (Bowen 2015) and provides a concise comparison of several methods. Of these, ovariectomy via colpotomy was found to be relatively safe when practiced by an experienced surgeon and was associated with the shortest duration of potential complications after the operation. The panel discussed the potential for evisceration through the vaginal incision with this procedure. In marked contrast to a suggestion by the NAS report (2013), this panel of veterinarians identified evisceration as not being a probable risk associated with ovariectomy via colpotomy and “none of the panel participants had had this occur nor had heard of it actually occurring” (Bowen 2015).

Most spay surgeries on mares have low morbidity and with the help of medications, pain and discomfort can be mitigated. Pain management is an important aspect of any ovariectomy (Rowland et al. 2018); according to surgical protocols that would be used, a long-lasting direct anesthetic would be applied to the ovarian pedicle, and systemic analgesics in the form of butorphanol and flunixin meglumine would be administered, as is compatible with accepted animal husbandry practices. In a study of the effects of bilateral ovariectomy via colpotomy on 23 mares, Hooper and others (1993) reported that postoperative problems were minimal (1 in 23, or 4%). Hooper et al. (1993) noted that four other mares were reported by owners as having some problems after surgery, but that evidence as to the role the surgery played in those subsequent problems was inconclusive. In contrast Röcken et al. (2011) noted a morbidity of 10.8% for mares that were ovariectomized via a flank laparoscopy. “Although 5 mares in our study had problems (repeated colic in 2 mares, signs of lumbar pain in 1 mare, signs of bilateral hind limb pain in 1 mare, and clinical signs of peritonitis in 1 mare) after surgery, evidence is inconclusive in each as to the role played by surgery” (Hooper et al. 1993). A recent study showed a 2.5% complication rate where one mare of 39 showed signs of moderate colic after laparoscopic ovariectomy (Devick 2018 personal communication).

Behavioral Effects of Mare Sterilization

No fertility control method exists that does not affect physiology or behavior of a mare (NAS 2013). Any action taken to alter the reproductive capacity of an individual has the potential to affect hormone production and therefore behavioral interactions and ultimately population dynamics in unforeseen ways (Ransom et al. 2014). The health and behavioral effects of spaying wild horse mares that live with other fertile and infertile wild horses has not been well documented, but the literature review below can be used to make reasonable inferences about their likely behaviors.

Horses are anovulatory (do not ovulate/express estrous behavior) during the short days of late fall and early winter, beginning to ovulate as days lengthen and then cycling roughly every 21 days during the warmer months, with about 5 days of estrus (Asa et al. 1979, Crowell-Davis 2007). Estrus in mares is shown by increased frequency of proceptive behaviors: approaching and following the stallion, urinating, presenting the rear end, clitoral winking, and raising the tail

towards the stallion (Asa et al. 1979, Crowell-Davis 2007). In most mammal species other than primates, estrus behavior is not shown during the anovulatory period, and reproductive behavior is considered extinguished following spaying (Hart and Eckstein 1997). However, mares may continue to demonstrate estrus behavior during the anovulatory period (Asa et al. 1980). Similarly, ovariectomized mares may also continue to exhibit estrous behavior (Scott and Kunze 1977, Kamm and Hendrickson 2007, Crabtree 2016), with one study finding that 30% of mares showed estrus signs at least once after surgery (Roessner et al 2015) and only 60 percent of ovariectomized mares cease estrous behavior following surgery (Loesch and Rodgerson 2003). Mares continue to show reproductive behavior following ovariectomy due to non-endocrine support of estrus behavior, specifically steroids from the adrenal cortex. Continuation of this behavior during the non-breeding season has the function of maintaining social cohesion within a horse group (Asa et al. 1980, Asa et al. 1984, NAS 2013). This may be a unique response of the horse (Bertin et al. 2013), as spaying usually greatly reduces female sexual behavior in companion animals (Hart and Eckstein 1997). In six ponies, mean monthly plasma luteinizing hormone levels in ovariectomized mares were similar to intact mares during the anestrus season, and during the breeding season were similar to levels in intact mares at mid-estrus (Garcia and Ginther 1976).

The likely effects of spaying on mares' social interactions and group membership can be inferred from available literature, even though wild horses have rarely been spayed and released back into the wild, resulting in few studies that have investigated their behavior in free-roaming populations. Wild horses and burros are instinctually herd-bound and this behavior is expected to continue. Overall, the BLM anticipates that some spayed mares may continue to exhibit estrus behavior which could foster band cohesion. If free-ranging ovariectomized mares show estrous behavior and occasionally allow copulation, interest of the stallion may be maintained, which could foster band cohesion (NAS 2013). This last statement could be validated by the observations of group associations on the Sheldon NWR where feral mares were ovariectomized via colpotomy and released back on to the range with untreated horses of both sexes (Collins and Kasbohm 2016). No data were collected on inter- or intra-band behavior (e.g. estrous display, increased tending by stallions, etc.), during multiple aerial surveys in years following treatment, all treated individuals appeared to maintain group associations, and there were no groups consisting only of treated males or only of treated females (Collins and Kasbohm 2016). In addition, of solitary animals documented during surveys, there were no observations of solitary treated females (Collins and Kasbohm 2016). These data help support the expectation that ovariectomized mares would not lose interest in or be cast out of the social dynamics of a wild horse herd. As noted by the NAS (2013), the ideal fertility control method would not eliminate sexual behavior or change social structure substantially.

A study conducted for 15 days in January 1978 (Asa et al. 1980), compared the sexual behavior in ovariectomized and seasonally anovulatory (intact) pony mares and found that there were no statistical differences between the two conditions for any measure of proceptivity or copulatory behavior, or days in estrus. This may explain why treated mares at Sheldon NWR continued to be accepted into harem bands; they may have been acting the same as a non-pregnant mare. Five to ten percent of pregnant mares exhibit estrous behavior (Crowell-Davis 2007). Although the physiological cause of this phenomenon is not fully understood (Crowell-Davis 2007), it is thought to be a bonding mechanism that assists in the maintenance of stable social groups of

horses year round (Ransom et al. 2014b). The complexity of social behaviors among free-roaming horses is not entirely centered on reproductive receptivity, and fertility control treatments that suppress the reproductive system and reproductive behaviors should contribute to minimal changes to social behavior (Ransom et al. 2014b, Collins and Kasbohm 2016).

BLM expects that wild horse harem structures would continue to exist under the proposed action because fertile mares, stallions, and their foals would continue to be a component of the herd. It is not expected that spaying a subset of mares would significantly change the social structure or herd demographics (age and sex ratios) of fertile wild horses.

‘Foal stealing,’ where a near-term pregnant mare steals a neonate foal from a weaker mare, is unlikely to be a common behavioral result of including spayed mares in a wild horse herd. McDonnell (2012) noted that “foal stealing is rarely observed in horses, except under crowded conditions and synchronization of foaling,” such as in horse feed lots. Those conditions are not likely in the wild, where pregnant mares will be widely distributed across the landscape, and where the expectation is that parturition dates would be distributed across the normal foaling season.

Indirect Effects of Mare sterilization

The free-roaming behavior of wild horses is not anticipated to be affected by mare sterilization, as the definition of free-roaming is the ability to move without restriction by fences or other barriers within a HMA (BLM H-4700-1, 2010) and there are no permanent physical barriers being proposed.

In domestic animals, spaying is often associated with weight gain and associated increase in body fat (Fettman et al 1997, Becket et al 2002, Jeusette et al. 2006, Belsito et al 2009, Reichler 2009, Camara et al. 2014). Spayed cats had a decrease in fasting metabolic rate, and spayed dogs had a decreased daily energy requirement, but both had increased appetite (O’Farrell & Peachey 1990, Hart and Eckstein 1997, Fettman et al. 1997, Jeusette et al. 2004). In wild horses, contracepted mares tend to be in better body condition than mares that are pregnant or that are nursing foals (Nuñez et al. 2010); the same improvement in body condition is likely to take place in spayed mares. In horses spaying has the potential to increase risk of equine metabolic syndrome (leading to obesity and laminitis), but both blood glucose and insulin levels were similar in mares before and after ovariectomy over the short-term (Bertin et al. 2013). In wild horses the quality and quantity of forage is unlikely to be sufficient to promote over-eating and obesity.

Coit et al. (2009) demonstrated that spayed dogs have elevated levels of LH-receptor and GnRH-receptor mRNA in the bladder tissue, and lower contractile strength of muscles. They noted that urinary incontinence occurs at elevated levels in spayed dogs and in post-menopausal women. Thus, it is reasonable to suppose that some ovariectomized mares could also suffer from elevated levels of urinary incontinence.

Sterilization had no effect on movements and space use of feral cats or brushtail possums (Ramsey 2007, Guttilla & Stapp 2010), or greyhound racing performance (Payne 2013). Rice field rats (*Rattus argentiventer*) tend to have a smaller home range in the breeding season, as they remain close to their litters to protect and nurse them. When surgically sterilized, rice field rats

had larger home ranges and moved further from their burrows than hormonally sterilized or fertile rats (Jacob et al. 2004). Spayed possums and foxes (*Vulpes vulpes*) had a similar core range area after spay surgery compared to before and were no more likely to shift their range than intact females (Saunders et al. 2002, Ramsey 2007).

The likely effects of spaying on mares' home range and habitat use can also be surmised from available literature. Bands of horses tend to have distinct home ranges, varying in size depending on the habitat and varying by season, but always including a water source, forage, and places where horses can shelter from inclement weather or insects (King and Gurnell 2005). It is unlikely that spayed mares will change their spatial ecology but being emancipated from constraints of lactation may mean they can spend more time away from water sources and increase their home range size. Lactating mares need to drink every day, but during the winter when snow can fulfill water needs or when not lactating, horses can traverse a wider area (Feist & McCullough 1976, Salter 1979). During multiple aerial surveys in years following the mare ovariectomy study at the Sheldon NWR, it was documented that all treated individuals appeared to maintain group associations, no groups consisted only of treated females, and none of the solitary animals observed were treated females (Collins and Kasbohm 2016). Because treated females maintained group associations, this indicates that their movement patterns and distances may be unchanged.

Spaying wild horses does not change their status as wild horses under the WFRHBA (as amended). In terms of whether spayed mares would continue to exhibit the free-roaming behavior that defines wild horses, BLM does expect that spayed mares would continue to roam unhindered. Wild horse movements may be motivated by a number of biological impulses, including the search for forage, water, and social companionship that is not of a sexual nature. As such, a spayed animal would still be expected to have a number of internal reasons for moving across a landscape and, therefore, exhibiting 'free-roaming' behavior. Despite marginal uncertainty about subtle aspects of potential changes in habitat preference, there is no expectation that spaying wild horses will cause them to lose their free-roaming nature.

In this sense, a spayed wild mare would be just as much 'wild' as defined by the WFRHBA as any fertile wild mare, even if her patterns of movement differ slightly. Congress specified that sterilization is an acceptable management action (16 USC §1333.b.1). Sterilization is not one of the clearly defined events that cause an animal to lose its status as a wild free-roaming horse (16 USC §1333.2.C.d). Any opinions based on a semantic and subjective definition of what constitutes a 'wild' horse are not legally binding for BLM, which must adhere to the legal definition of what constitutes a wild free-roaming horse, based on the WFRHBA (as amended). BLM is not obliged to base management decisions on personal opinions, which do not meet the BLM's principle and practice to "Use the best available scientific knowledge relevant to the problem or decision being addressed, relying on peer reviewed literature when it exists" (Kitchell et al. 2015).

Spaying is not expected to reduce mare survival rates on public rangelands. Individuals receiving fertility control often have reduced mortality and increased longevity due to being released from the costs of reproduction (Kirkpatrick and Turner 2008). Similar to contraception studies, in other wildlife species a common trend has been higher survival of sterilized females (Twigg et al. 2000, Saunders et al. 2002, Ramsey 2005, Jacob et al. 2008, Seidler and Gese 2012).

Observations from the Sheldon NWR provide some insight into long-term effects of ovariectomy on feral horse survival rates. The Sheldon NWR ovariectomized mares were returned to the range along with untreated mares. Between 2007 and 2014, mares were captured, a portion treated, and then recaptured. There was a minimum of 1 year between treatment and recapture; some mares were recaptured a year later, and some were recaptured several years later. The long-term survival rate of treated wild mares appears to be the same as that of untreated mares (Collins and Kasbohm 2016). Recapture rates for released mares were similar for treated mares and untreated mares.

Effects of Spaying on Bone Histology

The BLM knows of no scientific, peer-reviewed literature that documents bone density loss in mares following ovariectomy. A concern has been raised in an opinion article (Nock 2013) that ovary removal in mares could lead to bone density loss. That paper was not peer reviewed nor was it based on research in wild or domestic horses, so it does not meet the BLM's standard for "best available science" on which to base decisions (Kitchell et al. 2015). Hypotheses that are forwarded in Nock (2013) appear to be based on analogies from modern humans leading sedentary lives. Post-menopausal women have a greater chance of osteoporosis (Scholz-Ahrens et al. 1996), but BLM is not aware of any research examining bone loss in horses following ovariectomy. Bone loss in humans has been linked to reduced circulating estrogen. There have been conflicting results when researchers have attempted to test for an effect of reduced estrogen on animal bone loss rates in animal models; all experiments have been on laboratory animals, rather than free-ranging wild animals. While some studies found changes in bone cell activity after ovariectomy leading to decreased bone strength (Jerome et al. 1997, Baldock et al. 1998, Huang et al. 2002, Sigrist et al. 2007), others found that changes were moderate and transient or minimal (Scholz-Ahrens et al. 1996, Lundon et al. 1994, Zhang et al. 2007), and even returned to normal after 4 months (Sigrist et al. 2007).

Consistent and strenuous use of bones, for instance using jaw bones by eating hard feed, or using leg bones by travelling large distances, may limit the negative effects of estrogen deficiency on micro-architecture (Mavropoulos et al. 2014). The effect of exercise on bone strength in animals has been known for many years and has been shown experimentally (Rubin et al. 2001). Dr. Simon Turner, Professor Emeritus of the Small Ruminant Comparative Orthopaedic Laboratory at Colorado State University, conducted extensive bone density studies on ovariectomized sheep, as a model for human osteoporosis. During these studies, he did observe bone density loss on ovariectomized sheep, but those sheep were confined in captive conditions, fed twice a day, had shelter from inclement weather, and had very little distance to travel to get food and water (Simon Turner, Colorado State University Emeritus, written comm., 2015). Dr. Turner indicated that an estrogen deficiency (no ovaries) could potentially affect a horse's bone metabolism, just as it does in sheep and human females when they lead a sedentary lifestyle, but indicated that the constant weight bearing exercise, coupled with high exposure to sunlight ensuring high vitamin D levels, are expected to prevent bone density loss (Simon Turner, Colorado State University Emeritus, written comm., 2015).

Home range size of horses in the wild has been described as 4.2 to 30.2 square miles (Green and Green 1977) and 28.1 to 117 square miles (Miller 1983). A study of distances travelled by feral horses in "outback" Australia shows horses travelling between 5 and 17.5 miles per 24-hour

period (Hampson et al. 2010a), travelling about 11 miles a day even in a very large paddock (Hampson et al. 2010b). Thus, extensive movement patterns of wild horses are expected to help prevent bone loss. The expected daily movement distance would be far greater in the context of larger pastures typical of BLM long-term holding facilities in off-range pastures. A horse would have to stay on stall rest for years after removal of the ovaries in order to develop osteoporosis (Simon Turner, Colorado State University Emeritus, written comm., 2015) and that condition does not apply to any wild horses turned back to the range or any wild horses that go into off-range pastures.

G.4.3. Genetic Effects of Spaying and Neutering

It is true that spayed females and neutered males are unable to contribute to the genetic diversity of the herd. BLM is not obligated to ensure that any given individual in a herd has the chance to sire a foal and pass on genetic material. Management practices in the BLM Wild Horse and Burro Handbook (2010) include measures to increase population genetic diversity in reproducing herds where monitoring reveals a cause for concern about low levels of observed heterozygosity. These measures include increasing the sex ratio to a greater percentage of fertile males than fertile females (and thereby increasing the number of males siring foals) and bringing new animals into a herd from elsewhere.

In herds that are managed to be non-reproducing, it is not a concern to maintain genetic diversity because the management goal would be that animals in such a herd would not breed.

In reproducing herds where large numbers of wild horses have recent and / or an ongoing influx of breeding animals from other areas with wild or feral horses, spaying and neutering is not expected to cause an unacceptable loss of genetic diversity or an unacceptable increase in the inbreeding coefficient. In any diploid population, the loss of genetic diversity through inbreeding or drift can be prevented by large effective breeding population sizes (Wright 1931) or by introducing new potential breeding animals (Mills and Allendorf 1996). The NAS report (2013) recommended that single HMAs should not be considered as isolated genetic populations. Rather, managed herds of wild horses should be considered as components of interacting metapopulations, with the potential for interchange of individuals and genes taking place as a result of both natural and human-facilitated movements. It is worth noting that, although maintenance of genetic diversity at the scale of the overall population of wild horses is an intuitive management goal, there are no existing laws or policies that require BLM to maintain genetic diversity at the scale of the individual herd management area or complex. Also, there is no Bureau-wide policy that requires BLM to allow each female in a herd to reproduce before she is treated with contraceptives. Introducing 1-2 mares every generation (about every 10 years) is a standard management technique that can alleviate potential inbreeding concerns (BLM 2010). The NAS report (2013) recommended that managed herds of wild horses would be better viewed as components of interacting metapopulations, with the potential for interchange of individuals and genes taking place as a result of both natural and human-facilitated movements.

In the last 10 years, there has been a high realized growth rate of wild horses in most areas administered by the BLM. As a result, most alleles that are present in any given mare are likely to already be well represented in her siblings, cousins, and more distant relatives on the HMA. With the exception of horses in a small number of well-known HMAs that contain a relatively

high fraction of alleles associated with old Spanish horse breeds (NAS 2013), the genetic composition of wild horses in lands administered by the BLM is consistent with admixtures from domestic breeds. The NAS report (2013) includes information (pairwise genetic 'fixation index' values for sampled WH&B herds) confirming that WH&B in the vast majority of HMAs are genetically similar to animals in multiple other HMAs. As a result, in most HMAs, applying fertility control to a subset of mares is not expected to cause irreparable loss of genetic diversity. Improved longevity and an aging population are expected results of contraceptive treatment that can provide for lengthening generation time; this result would be expected to slow the rate of genetic diversity loss (Hailer et al. 2006). Based on a population model, Gross (2000) found that a strategy to preferentially treat young animals with a contraceptive led to more genetic diversity being retained than either a strategy that preferentially treats older animals, or a strategy with periodic gathers and removals.

Roelle and Oyler-McCance (2015) used the VORTEX population model to simulate how different rates of mare sterility would influence population persistence and genetic diversity, in populations with high or low starting levels of genetic diversity, various starting population sizes, and various annual population growth rates. Although those results are specific to mares, some inferences about potential effects of stallion sterilization may also be made from their results. Roelle and Oyler-McCance (2015) showed that the risk of the loss of genetic heterozygosity is extremely low except in cases where all of the following conditions are met: starting levels of genetic diversity are low, initial population size is 100 or less, the intrinsic population growth rate is low (5% per year), and very large fractions of the population are permanently sterilized. Given that 94 of 102 wild horse herds sampled for genetic diversity did not meet a threshold for concern (NAS 2013), the starting level of genetic diversity in most wild-horse herds is relatively high.

In a breeding herd where more than 85% of males in a population are sterile, there could be genetic consequences of reduced heterozygosity and increased inbreeding coefficients, as it would potentially allow a very small group of males to dominate the breeding (e.g., Saltz et al. 2000). Such genetic consequences could be mitigated by natural movements or human-facilitated translocations (BLM 2010). Garrott and Siniff's (1992) model predicts that gelding 50-80% of mature males in the population would result in reduced, but not halted, mare fertility rates. However, neutering males tends to have short-lived effects, because within a few years after any male sterilization treatment, a number of fertile male colts would become sexually mature stallions who could contribute genetically to the herd.

Literature Cited; Spaying and Neutering

- Angle, M., J. W. Turner Jr., R. M. Kenney, and V. K. Ganjam. 1979. Androgens in feral stallions. Pages 31–38 in Proceedings of the Symposium on the Ecology and Behaviour of Wild and Feral Equids, University of Wyoming, Laramie.
- Asa, C. S., D. A. Goldfoot, and O. J. Ginther. 1979. Sociosexual behavior and the ovulatory cycle of ponies (*Equus caballus*) observed in harem groups. *Hormones and Behavior* 13:49–65.

- Asa, C. S., D. A. Goldfoot, M. C. Garcia, and O. J. Ginther. 1980a. Dexamethasone suppression of sexual behavior in the ovariectomized mare. *Hormones and Behavior*.
- Asa, C., D. A. Goldfoot, M. C. Garcia, and O. J. Ginther. 1980b. Sexual behavior in ovariectomized and seasonally anovulatory pony mares (*Equus caballus*). *Hormones and Behavior* 14:46–54.
- Asa, C., D. Goldfoot, M. Garcia, and O. Ginther. 1984. The effect of estradiol and progesterone on the sexual behavior of ovariectomized mares. *Physiology and Behavior* 33:681–686.
- Asa, C. S. 1999. Male reproductive success in free-ranging feral horses. *Behavioural Ecology and Sociobiology* 47:89–93.
- Ashley, M.C., and D.W. Holcombe. 2001. Effects of stress induced by gathers and removals on reproductive success of feral horses. *Wildlife Society Bulletin* 29:248-254.
- Baldock, P. A. J., H. A. Morris, A. G. Need, R. J. Moore, and T. C. Durbridge. 1998. Variation in the short - term changes in bone cell activity in three regions of the distal femur immediately following ovariectomy. *Journal of Bone and Mineral Research* 13:1451–1457.
- Bartholow, J.M. 2004. An economic analysis of alternative fertility control and associated management techniques for three BLM wild horse herds. USGS Open-File Report 2004-1199.
- Beckett, T., A. E. Tchernof, and M. J. Toth. 2002. Effect of ovariectomy and estradiol replacement on skeletal muscle enzyme activity in female rats. *Metabolism* 51:1397–1401.
- Belsito, K. R., B. M. Vester, T. Keel, T. K. Graves, and K. S. Swanson. 2008. Impact of ovariohysterectomy and food intake on body composition, physical activity, and adipose gene expression in cats. *Journal of Animal Science* 87:594–602.
- Berger, J. 1986. *Wild horses of the Great Basin*. University of Chicago Press, Chicago.
- Bertin, F. R., K. S. Pader, T. B. Lescun, and J. E. Sojka-Kritchevsky. 2013. Short-term effect of ovariectomy on measures of insulin sensitivity and response to dexamethasone administration in horses. *American Journal of Veterinary Research* 74:1506–1513.
- Bigolin, S., D.J. Fagundes, H.C. Rivoire, A.T. Negrini Fagundes, A.L. Negrini Fagundes. 2009. Transcervical hysteroscopic sterilization using cyanoacrylate: a long-term experimental study on sheep. *The Journal of Obstetrics and Gynaecology Research* 35:1012-1018.
- Bowen, Z. 2015. Assessment of spay techniques for mare in field conditions. Letter from US Geological Survey Fort Collins Science Center to D. Bolstad, BLM. November 24, 2015. Appendix D in Bureau of Land Management, 2016, Mare Sterilization Research Environmental Assessment, DOI-BLM-O-B000-2015-055-EA, Hines, Oregon.

- BLM. 2010. BLM-4700-1 Wild Horses and Burros Management Handbook. Washington, D.C.
- BLM. 2011. Barren Valley Complex Wild Horse gather Plan. Final Environmental Assessment. DOI-BLM-OR-V040-2011-011-EA. BLM Oregon, Vale District / Jordan Field Office.
- BLM. 2012. Final Environmental Assessment Challis Wild Horse Gather Plan. DOI-BLM-ID-1030-2012-0006-EA. BLM Idaho, Challis Field Office.
- BLM. 2015. Instruction Memorandum 2015-151; Comprehensive animal welfare program for wild horse and burro gathers. Washington, D.C.
- BLM. 2016. Population Control Research Wild Horse Gather for the Conger and Frisco Herd Management Areas. Final Environmental Assessment. DOI-BLM-UT-W020-2015-0017-EA. BLM Utah, West Desert District.
- Borsberry, S. 1980. Libidinous behaviour in a gelding. *Veterinary Record* 106:89–90.
- Camara, C., L.-Y. Zhou, Y. Ma, L. Zhu, D. Yu, Y.-W. Zhao, and N.-H. Yang. 2014. Effect of ovariectomy on serum adiponectin levels and visceral fat in rats. *Journal of Huazhong University of Science and Technology [Medical Sciences]* 34:825–829.
- Chaudhuri, M., and J. R. Ginsberg. 1990. Urinary androgen concentrations and social status in two species of free ranging zebra (*Equus burchelli* and *E. grevyi*). *Reproduction* 88:127–133.
- Coit V. A., F. J. Dowell, and N. P. Evans. 2009. Neutering affects mRNA expression levels for the LH-and GnRH-receptors in the canine urinary bladder. *Theriogenology* 71:239-247.
- Colborn, D. R., D. L. Thompson, T. L. Roth, J. S. Capehart, and K. L. White. 1991. Responses of cortisol and prolactin to sexual excitement and stress in stallions and geldings. *Journal of Animal Science* 69:2556–2562.
- Collins, G. H., and J. W. Kasbohm. 2016. Population dynamics and fertility control of feral horses. *Journal of Wildlife Management* 81: 289-296.
- Costantini, R. M., J. H. Park, A. K. Beery, M. J. Paul, J. J. Ko, and I. Zucker. 2007. Post-castration retention of reproductive behavior and olfactory preferences in male Siberian hamsters: Role of prior experience. *Hormones and Behavior* 51:149–155.
- Crabtree, J. R. 2016. Can ovariectomy be justified on grounds of behaviour? *Equine Veterinary Education* 28: 58–59.
- Creel, S., B. Dantzer, W. Goymann, and D.R. Rubenstein. 2013. The ecology of stress: effects of the social environment. *Functional Ecology* 27:66-80.
- Crowell-Davis, S. L. 2007. Sexual behavior of mares.

- Deniston, R. H. 1979. The varying role of the male in feral horses. Pages 93–38 in Proceedings of the Symposium on the Ecology and Behaviour of Wild and Feral Equids, University of Wyoming, Laramie.
- de Seve, C.W. and S.L. Boyles-Griffin. 2013. An economic model demonstrating the long-term cost benefits of incorporating fertility control into wild horse (*Equus caballus*) management in the United States. *Journal of Zoo and Wildlife Medicine* 44(4s:S34-S37).
- Devick, I.F., B.S. Leise, S.Rao, and D.A. Hendrickson. 2018. Evaluation of post-operative pain after active desufflation at completion of laparoscopy in mares undergoing ovariectomy. *Canadian Veterinary Journal* 59:261-266.
- Dixson, A. F. 1993. Sexual and aggressive behaviour of adult male marmosets (*Callithrix jacchus*) castrated neonatally, prepubertally, or in adulthood. *Physiology and Behavior* 54:301–307.
- Dunbar, I. F. 1975. Behaviour of castrated animals. *The Veterinary Record* 92–93.
- Eagle, T. C., C. S. Asa, R. A. Garrott, E. D. Plotka, D. B. Siniff, and J. R. Tester. 1993. Efficacy of dominant male sterilization to reduce reproduction in feral horses. *Wildlife Society Bulletin* 21:116–121.
- Easley, J.T., K.C. McGilvray, D.A. Hendrickson, J. Bruemmer, and E.S. Hackett. 2018. Vessel sealer and divider instrument temperature during laparoscopic ovariectomy in horses. *Veterinary Surgery* 47: O26-O31.
- Evans, J. W., A. Borton, H. F. Hintz, and L. D. Van Vleck. 1977. *The Horse*. San Francisco, California: W.H. Freeman and Company. Pages 373–377.
- Feh, C. 1999. Alliances and reproductive success in Camargue stallions. *Animal Behaviour* 57:705–713.
- Feist, J. D., and D. R. McCullough. 1976. Behavior patterns and communication in feral horses. *Zeitschrift für Tierpsychologie* 41:337–371.
- Feist, J. D., and D.R. McCullough. 1976. Behavior patterns and communication in feral horses. *Zeitschrift für Tierpsychologie* 41:337–371.
- Fettman, M. J., C. A. Stanton, L. L. Banks, D. W. Hamar, D. E. Johnson, R. L. Hegstad, and S. Johnston. 1997. Effects of neutering on bodyweight, metabolic rate and glucose tolerance of domestic cats. *Research in Veterinary Science* 62:131–136.
- Fonner, R. and A.K. Bohara. 2017. Optimal control of wild horse populations with nonlethal methods. *Land Economics* 93:390-412.
- Garcia, M. C., and O. J. Ginther. 1976. Effects of Ovariectomy and Season on Plasma Luteinizing Hormone in Mares. *Endocrinology* 98:958–962.

- Garrott, R.A., and D.B. Siniff. 1992. Limitations of male-oriented contraception for controlling feral horse populations. *Journal of Wildlife Management* 56:456-464.
- Garrott, R.A., and M.K. Oli. 2013. A Critical Crossroad for BLM's Wild Horse Program. *Science* 341:847-848.
- Getman, L.M. 2009. Review of castration complications: strategies for treatment in the field. *AAEP Proceedings* 55:374-378.
- Green, N.F. and H.D. Green. 1977. The wild horse population of Stone Cabin Valley Nevada: a preliminary report. In *Proceedings, National Wild Horse Forum*. University of Nevada Reno Cooperative Extension Service.
- Gross, J.E. 2000. A dynamic simulation model for evaluating effects of removal and contraception on genetic variation and demography of Pryor Mountain wild horses. *Biological Conservation* 96:319-330.
- Gutilla, D. A., and P. Stapp. 2010. Effects of sterilization on movements of feral cats at a wildland–urban interface. *Journal of Mammalogy* 91:482–489.
- Hailer, F., B. Helander, A.O. Folkestad, S.A. Ganusevich, S. Garstad, P. Hauff, C. Koren, T. Nygård, V. Volke, C. Vilà, and H. Ellegren. 2006. Bottlenecked but long-lived: high genetic diversity retained in white-tailed eagles upon recovery from population decline. *Biology Letters* 2:316-319.
- Hampson, B. A., M. A. De Laat, P. C. Mills, and C. C. Pollitt. 2010a. Distances travelled by feral horses in ‘outback’ Australia. *Equine Veterinary Journal, Suppl.* 38:582–586.
- Hampson, B. A., J. M. Morton, P. C. Mills, M. G. Trotter, D. W. Lamb, and C. C. Pollitt. 2010b. Monitoring distances travelled by horses using GPS tracking collars. *Australian Veterinary Journal* 88:176–181.
- Hampton, J.O., T.H. Hyndman, A. Barnes, and T. Collins. 2015. Is wildlife fertility control always humane? *Animals* 5:1047-1071.
- Hart, B. L. 1968. Role of prior experience in the effects of castration on sexual behavior of male dogs. *Journal of Comparative and Physiological Psychology* 66:719–725.
- Hart, B. L., and T. O. A. C. Jones. 1975. Effects of castration on sexual behavior of tropical male goats. *Hormones and Behavior* 6:247–258.
- Hart, B. L., and R. A. Eckstein. 1997. The role of gonadal hormones in the occurrence of objectionable behaviours in dogs and cats. *Applied Animal Behaviour Science* 52:331–344.

- Hobbs, N.T., D.C. Bowden and D.L. Baker. 2000. Effects of Fertility Control on Populations of Ungulates: General, Stage-Structured Models. *Journal of Wildlife Management* 64:473-491.
- Holtan, D. W., E. L. Squires, D. R. Lapin, and O. J. Ginther. 1979. Effect of ovariectomy on pregnancy in mares. *Journal of Reproduction and Fertility, Supplement* 27:457–463.
- Hooper, R. N., T. S. Taylor, D. D. Varner, and B. T. L. 1993. Effects of bilateral ovariectomy via colotomy in mares: 23 cases (1984-1990). *Journal of the American Veterinary Medical Association* 203:1043–1046.
- Huang, R. Y., L. M. Miller, C. S. Carlson, and M. R. Chance. 2002. Characterization of bone mineral composition in the proximal tibia of *Cynomolgus* monkeys: effect of ovariectomy and nandrolone decanoate treatment. *Bone* 30:492–497.
- Jacob, J., G. R. Singleton, and L. A. Hinds. 2008. Fertility control of rodent pests. *Wildlife Research* 35:487.
- Jerome, C. P., C. H. Turner, and C. J. Lees. 1997. Decreased bone mass and strength in ovariectomized cynomolgus monkeys (*Macaca fascicularis*). *Calcified Tissue International* 60:265–270.
- Jeusette, I., J. Detilleux, C. Cuvelier, L. Istasse, and M. Diez. 2004. Ad libitum feeding following ovariectomy in female Beagle dogs: effect on maintenance energy requirement and on blood metabolites. *Journal of Animal Physiology and Animal Nutrition* 88:117–121.
- Jeusette, I., S. Daminet, P. Nguyen, H. Shibata, M. Saito, T. Honjoh, L. Istasse, and M. Diez. 2006. Effect of ovariectomy and ad libitum feeding on body composition, thyroid status, ghrelin and leptin plasma concentrations in female dogs. *Journal of Animal Physiology and Animal Nutrition* 90:12–18.
- Jewell, P. A. 1997. Survival and behaviour of castrated Soay sheep (*Ovis aries*) in a feral island population on Hirta, St. Kilda, Scotland. *Journal of Zoology* 243:623–636.
- Kamm, J. L., and D. A. Hendrickson. 2007. Clients' perspectives on the effects of laparoscopic ovariectomy on equine behavior and medical problems. *Journal of Equine Veterinary Science* 27:435–438.
- Kaseda, Y., H. Ogawa, and A. M. Khalil. 1997. Causes of natal dispersal and emigration and their effects on harem formation in Misaki feral horses. *Equine Veterinary Journal* 29:262–266.
- Khalil, A.M., N. Murakami, and Y. Kaseda. 1998. Relationship between plasma testosterone concentrations and age, breeding season, and harem size in Misaki feral horses. *Journal of Veterinary Medical Science* 60:643-645.

- Khalil, A. M., and N. Murakami. 1999. Effect of natal dispersal on the reproductive strategies of the young Misaki feral stallions. *Applied Animal Behaviour Science* 62:281–291.
- King, S.R.B., and J. Gurnell. 2005. Habitat use and spatial dynamics of takhi introduced to Hustai National Park, Mongolia. *Biological Conservation* 124:277–290.
- King, S.R.B., and J. Gurnell. 2006. Scent-marking behaviour by stallions: an assessment of function in a reintroduced population of Przewalski horses (*Equus ferus przewalskii*). *Journal of Zoology* 272:30–36.
- Kirkpatrick, J. 2012. Sworn statement of Dr. Jay Kirkpatrick. Unpublished record of opinion.
- Kirkpatrick, J. F., and A. Turner. 2008. Achieving population goals in a long-lived wildlife species (*Equus caballus*) with contraception. *Wildlife Research* 35:513.
- Kitchell, K., S. Cohn, R. Falise, H. Hadley, M. Herder, K. Libby, K. Muller, T. Murphy, M. Preston, M.J. Rugwell, and S. Schlanger. 2015. Advancing science in the BLM: an implementation strategy. Department of the Interior, BLM, Washington DC.
- Lee, M., and D. A. Hendrickson. 2008. A review of equine standing laparoscopic ovariectomy. *Journal of Equine Veterinary Science* 28:105–111.
- Line, S. W., B. L. Hart, and L. Sanders. 1985. Effect of prepubertal versus postpubertal castration on sexual and aggressive behavior in male horses. *Journal of the American Veterinary Medical Association* 186:249–251.
- Linklater, W. L., and E. Z. Cameron. 2000. Distinguishing cooperation from cohabitation: the feral horse case study. *Animal Behaviour* 59:F17–F21.
- Loesch, D. A., and D. H. Rodgerson. 2003. Surgical approaches to ovariectomy in mares. *Continuing Education for Veterinarians* 25:862–871.
- Lundon, K., M. Dumitriu, and M. Gryn timer. 1994. The long-term effect of ovariectomy on the quality and quantity of cancellous bone in young macaques. *Bone and Mineral* 24:135–149.
- Mavropoulos, A., S. Kiliaridis, R. Rizzoli, and P. Ammann. 2014. Normal masticatory function partially protects the rat mandibular bone from estrogen-deficiency induced osteoporosis. *Journal of Biomechanics* 47:2666–2671.
- McDonnell, S.M. 2012. Mare and foal behavior. *American Association of Equine Practitioners Proceedings* 58:407–410.
- McKinnon, A.O., and J.R. Vasey. 2007. Selected reproductive surgery of the broodmare. Pages 146–160 in *Current therapy in equine reproduction*, J.C. Samper, J.F. Pycoc, and A.O. McKinnon, eds. Saunders Elsevier, St. Louis, Missouri.

- Miller, R. 1983. Seasonal Movements and Home Ranges of Feral Horse Bands in Wyoming's Red Desert. *Journal of Range Management* 36:199–201.
- Mills, L.S. and F.W. Allendorf. 1996. The one - migrant - per - generation rule in conservation and management. *Conservation Biology* 10:1509-1518.
- National Research Council of the National Academies of Sciences (NAS). 2013. Using science to improve the BLM wild horse and burro program: a way forward. National Academies Press. Washington, DC.
- National Research Council of the National Academies of Sciences (NAS). 2015. Review of proposals to the Bureau of Land Management on Wild Horse and Burro sterilization or contraception, a letter report. Committee for the review of proposals to the Bureau of Land Management on Wild Horse and Burro Sterilization or Contraception. Appendix B in: BLM, 2016, Mare sterilization research Environmental Assessment DOI-BLM-OR-B000-2015-0055-EA, BLM Burns District Office, Hines, Oregon.
- Nelson, K. J. 1980. Sterilization of dominant males will not limit feral horse populations. USDA Forest Service Research Paper RM-226.
- Nickolmann, S., S. Hoy, and M. Gauly. 2008. Effects of castration on the behaviour of male llamas (*Lama glama*). *Tierärztliche Praxis Großtiere* 36:319–323.
- Nock, B. 2013. Liberated horsemanship: menopause...and wild horse management. Warrenton, Missouri: Liberated Horsemanship Press.
- Nock, B. 2017. Gelding is likely to cause wild horses undo suffering. Unpublished record of opinion.
- Núñez, C.M., J.S. Adelman, and D.I. Rubenstein. 2010. Immunocontraception in wild horses (*Equus caballus*) extends reproductive cycling beyond the normal breeding season. *PLoS one*, 5(10), p.e13635.
- Núñez, C.M., J.S. Adelman, H.A. Carr, C.M. Alvarez, and D.I. Rubenstein. 2017. Lingering effects of contraception management on feral mare (*Equus caballus*) fertility and social behavior. *Conservation Physiology* 5(1): cox018; doi:10.1093/conphys/cox018.
- O'Farrell, V., and E. Peachey. 1990. Behavioural effects of ovariectomy on bitches. *Journal of Small Animal Practice* 31:595–598.
- Pader, K., L. J. Freeman, P. D. Constable, C. C. Wu, P. W. Snyder, and T. B. Lescun. 2011. Comparison of Transvaginal Natural Orifice Transluminal Endoscopic Surgery (NOTES®) and Laparoscopy for Elective Bilateral Ovariectomy in Standing Mares. *Veterinary Surgery* 40:998–1008.
- Payne, R. M. 2013. The effect of spaying on the racing performance of female greyhounds. *The Veterinary Journal* 198:372–375.

- Pearce, O. 1980. Libidinous behaviour in a gelding. *Veterinary Record* 106:207–207.
- Prado, T., and J. Schumacher. 2017. How to perform ovariectomy through a colpotomy. *Equine Veterinary Education* 13: doi: 10.1111/eve.12801
- Ramsey, D. 2005. Population dynamics of brushtail possums subject to fertility control. *Journal of Applied Ecology* 42:348–360.
- Ramsey, D. 2007. Effects of fertility control on behavior and disease transmission in brushtail possums. *Journal of Wildlife Management* 71:109–116.
- Ransom, J. I., and B. S. Cade. 2009. Quantifying Equid Behavior--A Research Ethogram for Free-Roaming Feral Horses. Publications of the US Geological Survey. U.S. Geological Survey Techniques and Methods 2-A9.
- Ransom, J.I., J.G. Powers, N.T. Hobbs, and D.L. Baker. 2014a. Ecological feedbacks can reduce population-level efficacy of wildlife fertility control. *Journal of Applied Ecology* 51:259-269.
- Ransom, J.I., J.G. Powers, H.M. Garbe, M.W. Oehler, T.M. Nett, and D.L. Baker. 2014b. Behavior of feral horses in response to culling and GnRH immunocontraception. *Applied Animal Behaviour Science* 157: 81-92.
- Reichler, I. M. 2009. Gonadectomy in Cats and Dogs: A Review of Risks and Benefits. *Reproduction in Domestic Animals* 44:29–35.
- Rios, J. F. I., and K. Houpt. 1995. Sexual behavior in geldings. *Applied Animal Behaviour Science* 46:133–133.
- Roelle, J. E., F. J. Singer, L. C. Zeigenfuss, J. I. Ransom, L. Coates-Markle, and K. A. Schoenecker. 2010. Demography of the Pryor Mountain Wild Horses, 1993–2007. pubs.usgs.gov. U.S. Geological Survey Scientific Investigations Report 2010-5125.
- Röcken, M., G. Mosel, K. Seyrek-Intas, D. Seyrek-Intas, F. Litzke, J. Verver, and A. B. M. Rijkenhuizen. 2011. Unilateral and Bilateral Laparoscopic Ovariectomy in 157 Mares: A Retrospective Multicenter Study. *Veterinary Surgery* 40:1009–1014.
- Roelle, J.E. and S.J. Oyler-McCance. 2015. Potential demographic and genetic effects of a sterilant applied to wild horse mares. US Geological Survey Open-file Report 2015-1045.
- Roessner, H. A., K.A. Kurtz, and J.P. Caron. 2015. Laparoscopic ovariectomy diminishes estrus-associated behavioral problems in mares. *Journal of Equine Veterinary Science* 35: 250–253 (2015).
- Rowland, A.L., K.G. Glass, S.T. Grady, K.J. Cummings, K. Hinrichs, and A.E. Watts. 2018. Influence of caudal epidural analgesia on cortisol concentrations and pain-related

- behavioral responses in mares during and after ovariectomy via colpotomy. *Veterinary Surgery* 2018:1-7. DOI: 10.1111/vsu.12908
- Rubin, C., A. S. Turner, S. Bain, C. Mallinckrodt, and K. McLeod. 2001. Low mechanical signals strengthen long bones. *Nature* 412:603–604.
- Rutberg, A. 2011. Re: Modified decision record, WY-040-EA11-124. Unpublished record of opinion.
- Salter, R. E. Biogeography and habitat-use behavior of feral horses in western and northern Canada. In *Symposium on the Ecology and Behaviour of Wild and Feral Equids* 129–141 (1979).
- Saltz, D., M. Rowen, and D. I. Rubenstein. 2000. The effect of space - use patterns of reintroduced Asiatic wild ass on effective population size. *Conservation Biology* 14:1852–1861.
- Saunders, G., J. McIlroy, M. Berghout, B. Kay, E. Gifford, R. Perry, and R. van de Ven. 2002. The effects of induced sterility on the territorial behaviour and survival of foxes. *Journal of Applied Ecology* 39:56–66.
- Scholz-Ahrens, K. E., G. Delling, P. W. Jungblut, E. Kallweit, and C. A. Barth. 1996. Effect of ovariectomy on bone histology and plasma parameters of bone metabolism in nulliparous and multiparous sows. *Zeitschrift für Ernährungswissenschaft* 35:13–21.
- Schumacher, J. 1996. Complications of castration. *Equine Veterinary Education* 8:254-259.
- Schumacher, J. 2006. Why do some castrated horses still act like stallions, and what can be done about it? *Compendium Equine Edition Fall*: 142–146.
- Scott, E. A., and D. J. Kunze. 1977. Ovariectomy in the mare: presurgical and postsurgical considerations. *The Journal of Equine Medicine and Surgery* 1:5–12.
- Searle, D., A.J. Dart, C.M. Dart, and D.R. Hodgson. 1999. Equine castration: review of anatomy, approaches, techniques and complications in normal, cryptorchid and monorchid horses. *Australian Veterinary Journal* 77:428-434.
- Seidler, R. G., and E. M. Gese. 2012. Territory fidelity, space use, and survival rates of wild coyotes following surgical sterilization. *Journal of Ethology* 30:345–354.
- Shoemaker, R., Bailey, J., Janzen, E. and Wilson, D.G., 2004. Routine castration in 568 draught colts: incidence of evisceration and omental herniation. *Equine Veterinary Journal*, 36:336-340.
- Shoemaker, R. W., E. K. Read, T. Duke, and D. G. Wilson. 2004. In situ coagulation and transection of the ovarian pedicle: an alternative to laparoscopic ovariectomy in juvenile horses. *Canadian Journal of Veterinary Research* 68:27-32.

- Sigrist, I. M., C. Gerhardt, M. Alini, E. Schneider, and M. Egermann. 2007. The long-term effects of ovariectomy on bone metabolism in sheep. *Journal of Bone and Mineral Metabolism* 25:28–35.
- Sigurjónsdóttir, H., M. C. Van Dierendonck, S. Snorrason, and A. G. Thorhallsdóttir. 2003. Social relationships in a group of horses without a mature stallion. *Behaviour* 140:783–804.
- Smith, J. A. 1974. Proceedings: Masculine behaviour in geldings. *The Veterinary Record* 94:160–160.
- Thompson, D. L., Jr, B. W. Pickett, E. L. Squires, and T. M. Nett. 1980. Sexual behavior, seminal pH and accessory sex gland weights in geldings administered testosterone and (or) estradiol-17. *Journal of Animal Science* 51:1358–1366.
- Twigg, L. E., T. J. Lowe, G. R. Martin, A. G. Wheeler, G. S. Gray, S. L. Griffin, C. M. O'Reilly, D. J. Robinson, and P. H. Hubach. 2000. Effects of surgically imposed sterility on free-ranging rabbit populations. *Journal of Applied Ecology* 37:16–39.
- Tyler, S. 1972. The behaviour and social organisation of the New Forest ponies. *Animal Behaviour Monographs* 5:85–196.
- US Fish and Wildlife Service (USFWS). 2015. Endangered and Threatened Wildlife and Plants; 90-day findings on 31 petitions. *Federal Register* 80 (126):37568-37579.
- Van Dierendonck, M. C., H. De Vries, and M. B. H. Schilder. 1995. An analysis of dominance, its behavioural parameters and possible determinants in a herd of Icelandic horses in captivity. *Journal of Zoology* 45:362–385.
- Van Dierendonck, M. C., H. Sigurjónsdóttir, B. Colenbrander, and A. G. Thorhallsdóttir. 2004. Differences in social behaviour between late pregnant, post-partum and barren mares in a herd of Icelandic horses. *Applied Animal Behaviour Science* 89:283–297.
- Van Dierendonck, M. C., H. De Vries, M. B. H. Schilder, B. Colenbrander, A. G. Þorhallsdóttir, and H. Sigurjónsdóttir. 2009. Interventions in social behaviour in a herd of mares and geldings. *Applied Animal Behaviour Science* 116:67–73.
- Vinke, C. M., R. van Deijk, B. B. Houx, and N. J. Schoemaker. 2008. The effects of surgical and chemical castration on intermale aggression, sexual behaviour and play behaviour in the male ferret (*Mustela putorius furo*). *Applied Animal Behaviour Science* 115:104–121.
- Webley, G. E., and E. Johnson. 1982. Effect of ovariectomy on the course of gestation in the grey squirrel (*Sciurus carolinensis*). *Journal of Endocrinology* 93:423–426.
- Wright, S. 1931. Evolution in Mendelian populations. *Genetics* 16:97-159

Zhang, Y., W.-P. Lai, P.-C. Leung, C.-F. Wu, and M.-S. Wong. 2007. Short- to Mid-Term Effects of Ovariectomy on Bone Turnover, Bone Mass and Bone Strength in Rats. *Biological and Pharmaceutical Bulletin* 30:898–903.

G.5. Intrauterine Devices (IUDs)

Up through the present time (September 2020), BLM has not used intrauterine devices (IUDs) to control fertility as a wild horse and burro fertility control method on the range. The BLM has supported and continues to support research into the development and testing of effective and safe IUDs for use in wild horse mares (Baldrighi et al. 2017, Holyoak et al. unpublished data). However, existing literature on the use of IUDs in horses allows for inferences about expected effects of any management alternatives that might include use of IUDs, and support the apparent safety and efficacy of some types of IUDs for use in horses. Overall, as with other methods of population growth suppression, use of IUDs and other fertility control measures are expected to help reduce population growth rates, extend the time interval between gathers, and reduce the total number of excess animals that will need to be removed from the range.

The 2013 National Academies of Sciences (NAS) report considered IUDs and suggested that research should test whether IUDs cause uterine inflammation, and should also test how well IUDs stay in mares that live and breed with fertile stallions. Since that report, a recent study by Holyoak et al. (unpublished data) indicate that a flexible, inert, y-shaped, medical-grade silicone IUD design prevented pregnancies in all the domestic mares that retained the device, even when exposed to fertile stallions. Domestic mares in that study lived in large pastures, mating with fertile stallions. Biweekly ultrasound examinations showed that IUDs stayed in 75% of treated mares over the course of two breeding seasons. The IUDs were then removed so the researchers could monitor the mares' return to fertility. Uterine health, as measured in terms of inflammation, was not seriously affected by the IUDs, and most mares became pregnant within months after IUD removal. The overall results are consistent with results from an earlier study (Daels and Hughes 1995), which used O-shaped silicone IUDs.

IUDs are considered a temporary fertility control method that does not generally cause future sterility (Daels and Hughes 1995). Use of IUDs is an effective fertility control method in women, and IUDs have historically been used in livestock management, including in domestic horses. Insertion of an IUD can be a very rapid procedure, but it does require the mare to be temporarily restrained, such as in a squeeze chute. IUDs in mares may cause physiological effects including discomfort, infection, perforation of the uterus if the IUD is hard and angular, endometritis, uterine edema (Killian et al. 2008), and pyometra (Klabnik-Bradford et al. 2013). In women, deaths attributable to IUD use may be as low as 1.06 per million (Daels and Hughes 1995).

The exact mechanism by which IUDs prevent pregnancy is uncertain (Daels and Hughes 1995), but the presence of an IUD in the uterus may, like a pregnancy, prevent the mare from coming back into estrus (Turner et al. 2015). However, some domestic mares did exhibit repeated estrus cycles during the time when they had IUDs (Killian et al. 2008, Gradil et al. 2019). The main cause for an IUD to not be effective at contraception is its failure to stay in the uterus (Daels and Hughes 1995). As a result, one of the major challenges to using IUDs to control fertility in mares on the range is preventing the IUD from being dislodged or otherwise ejected over the course of daily activities, which could include, at times, frequent breeding.

At this time, it is thought that any IUD inserted into a pregnant mare may cause the pregnancy to terminate, which may also cause the IUD to be expelled. For that reason, it is expected that IUDs would only be inserted in non-pregnant (open) mares. Wild mares receiving IUDs would be checked for pregnancy prior to insertion of an IUD. This can be accomplished by transrectal palpation and/or ultrasound performed by a veterinarian. Pregnant mares would not receive an IUD. The IUD is inserted into the uterus using a thin, tubular applicator similar to a shielded culture tube, and would be inserted in a manner similar to that routinely used to obtain uterine cultures in domestic mares. If a mare has a zygote or very small, early phase embryo, it is possible that it will fail to be detected in screening, and may develop further, but without causing the expulsion of the IUD. Wild mares with IUDs would be individually marked and identified, so that they can be monitored occasionally and examined, if necessary, in the future, consistent with other BLM management activities.

Using metallic or glass marbles as IUDs may prevent pregnancy in horses (Nie et al. 2003) but can pose health risks to domestic mares (Turner et al. 2015, Freeman and Lyle 2015). Marbles may break into shards (Turner et al. 2015), and uterine irritation that results from marble IUDs may cause chronic, intermittent colic (Freeman and Lyle 2015). Metallic IUDs may cause severe infection (Klabnik-Bradford et al. 2013).

In domestic ponies, Killian et al. (2008) explored the use of three different IUD configurations, including a silastic polymer O-ring with copper clamps, and the “380 Copper T” and “GyneFix” IUDs designed for women. The longest retention time for the three IUD models was seen in the “T” device, which stayed in the uterus of several mares for 3-5 years. Reported contraception rates for IUD-treated mares were 80%, 29%, 14%, and 0% in years 1-4, respectively. They surmised that pregnancy resulted after IUD fell out of the uterus. Killian et al. (2008) reported high levels of progesterone in non-pregnant, IUD-treated ponies.

Soft IUDs may cause relatively less discomfort than hard IUDs (Daels and Hughes 1995). Daels and Hughes (1995) tested the use of a flexible O-ring IUD, made of silastic, surgical-grade polymer, measuring 40 mm in diameter; in five of six breeding domestic mares tested, the IUD was reported to have stayed in the mare for at least 10 months. In mares with IUDs, Daels and Hughes (1995) reported some level of uterine irritation but surmised that the level of irritation was not enough to interfere with a return to fertility after IUD removal.

More recently, several types of IUDs have been tested for use in breeding mares. When researchers attempted to replicate the O-ring study (Daels and Hughes 1995) in an USGS / Oklahoma State University (OSU) study with breeding domestic mares, using various configurations of silicone O-ring IUDs, the IUDs fell out at unacceptably high rates over time scales of less than 2 months (Baldrighi et al. 2017). Subsequently, the USGS / OSU researchers tested a Y-shaped IUD to determine retention rates and assess effects on uterine health; retention rates were greater than 75% for an 18-month period, and mares returned to good uterine health and reproductive capacity after removal of the IUDs (Holyoak et al., unpublished results). The University of Massachusetts has developed a magnetic IUD that has been effective at preventing estrus in non-breeding domestic mares (Gradil et al. 2019). After insertion in the uterus, the three subunits of the device are held together by magnetic forces as a flexible triangle. A metal detector can be used to determine whether the device is still present in the mare. In an early trial, two sizes of those magnetic IUDs fell out of breeding domestic mares at high rates (Holyoak et

al., unpublished results). In 2019, the magnetic IUD was used in two trials where mares were exposed to stallions, and in one where mares were artificially inseminated; in all cases, the IUDs were reported to stay in the mares without any pregnancy (Gradil 2019).

Literature Cited: Intrauterine Devices (IUDs)

- Baldrighi, J.M., C.C. Lyman, K. Hornberger, S.S. Germaine, A. Kane, and G.R. Holyoak. 2017. Evaluating the efficacy and safety of silicone O-ring intrauterine devices as a horse contraceptive through a captive breeding trial. *Clinical Theriogenology* 9:471.
- Daels, P.F, and J.P. Hughes. 1995. Fertility control using intrauterine devices: an alternative for population control in wild horses. *Theriogenology* 44:629-639.
- Freeman, C.E., and S.K. Lyle. 2015. Chronic intermittent colic in a mare attributed to uterine marbles. *Equine Veterinary Education* 27:469-473.
- Gradil, C. 2019. The Upod IUD: a potential simple, safe solution for long-term, reversible fertility control in feral equids. Oral presentation at the Free Roaming Equids and Ecosystem Sustainability Summit, Reno, Nevada.
- Gradil, C.M., C.K. Uricchio, and A. Schwarz. 2019. Self-Assembling Intrauterine Device (Upod) Modulation of the Reproductive Cycle in Mares. *Journal of Equine Veterinary Science* 83: 102690.
- Holyoak, G.R., C.C. Lyman, S. Wang, S.S. Germaine, C.O. Anderson, J.M. Baldrighi, N. Vemula, G.B. Rexabek, and A.J. Kane. Unpublished. Efficacy of a Y-design intrauterine device as a horse contraceptive. In review.
- Killian, G., D. Thain, N.K. Diehl, J. Rhyon, and L. Miller. 2008. Four-year contraception rates of mares treated with single-injection porcine zona pellucida and GnRH vaccines and intrauterine devices. *Wildlife Research* 35:531-539.
- Klabnik-Bradford, J., M.S. Ferrer, C. Blevins, and L. Beard. 2013. Marble-induced pyometra in an Appaloosa mare. *Clinical Theriogenology* 5: 410.
- Nie, G.J., K.E., Johnson, T.D. Braden, and J. G.W. Wenzel. 2003. Use of an intra-uterine glass ball protocol to extend luteal function in mares. *Journal of Equine Veterinary Science* 23:266-273.
- Turner, R.M., D.K. Vanderwall, and R. Stawecki. 2015. Complications associated with the presence of two intrauterine glass balls used for oestrus suppression in a mare. *Equine Veterinary Education* 27:340-343.